

Anti-EID3 antibody ab124447

[1 References](#) [2 图像](#)

概述

产品名称	Anti-EID3抗体
描述	兔多克隆抗体to EID3
宿主	Rabbit
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human 预测可用于: Mouse, Rat 
免疫原	Recombinant fragment, corresponding to amino acids 139-333 of Human EID3 (BC027612).
阳性对照	Human fetal brain, liver, kidney lysates, HeLa, K562 and HepG2 cell lysates; fetasl testis tissue.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Lyophilized:Reconstitute in 200ul sterile H2O.
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.02% Sodium azide Constituent: 1% BSA
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab124447于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/2000. Predicted molecular weight: 38 kDa.
IHC-P		1/100 - 1/500.

靶标

功能	Act as a repressor of nuclear receptor-dependent transcription possibly by interfering with CREBBP-dependent coactivation. May function as a coinhibitor of other CREBBP/EP300-dependent transcription factors.
组织特异性	Highly expressed in testis.
序列相似性	Belongs to the NSE4 family.
细胞定位	Nucleus. Cytoplasm. May shuttle between nucleus and cytoplasm.

图片



All lanes : Anti-EID3 antibody (ab124447) at 1/1000 dilution

Lane 1 : Human fetal brain lysate

Lane 2 : Human fetal liver lysate


Lane 3 : Human fetal kidney lysate

Lane 4 : HeLa cell lysate

Lane 5 : K562 cell lysate

Lane 6 : HepG2 cell lysate

Predicted band size: 38 kDa



ab124447 at 1/100 dilution staining EID3 in Human fetal testis tissue by Immunohistochemistry Formalin-fixed, Paraffin-embedded tissue.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors