

Product datasheet

Anti-ERG antibody [EPR3864] ab92513

重组 RabMAb


★★★★★ [17 Abreviews](#) [201 References](#) [17 图像](#)

概述

产品名称	Anti-ERG抗体[EPR3864]
描述	兔单克隆抗体[EPR3864] to ERG
宿主	Rabbit
特异性	This antibody also detects Fli-1.
经测试应用	适用于: WB, IHC-P, ICC/IF, Flow Cyt (Intra)
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Jurkat, HeLa and RAW 264.7 cell lysates; Rat brain and heart lysates. IHC-P: Human kidney, brain and prostate adenocarcinoma tissues; Fus A5 transgenic mouse prostate tissue; Mouse brain tissue. ICC/IF: Circulating tumor cells (CTCs) from a castrate-resistant prostate cancer (CRPC) patient; THP-1 cells. Flow Cyt (intra): THP-1 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K_D)	K _D = 8.90 x 10 ⁻¹⁰ M



[Learn more about K_D](#)

存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR3864
同种型	IgG

应用


The Abpromise guarantee **Abpromise™** 承诺保证使用 ab92513 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (1)	1/1000 - 1/10000. Predicted molecular weight: 55 kDa.
IHC-P	★★★★★ (3)	1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified, use 1/100 - 1/250.
ICC/IF	★★★★★ (3)	Use a concentration of 1 µg/ml. This product gave a positive signal in THP-1 (-ve: HCT116) fixed with 4% formaldehyde (10 min).
Flow Cyt (Intra)		Use at an assay dependent concentration.


靶标

功能	Transcriptional regulator. May participate in transcriptional regulation through the recruitment of SETDB1 histone methyltransferase and subsequent modification of local chromatin structure.
疾病相关	Defects in ERG are a cause of Ewing sarcoma (ES) [MIM:612219]. A highly malignant, metastatic, primitive small round cell tumor of bone and soft tissue that affects children and adolescents. It belongs to the Ewing sarcoma family of tumors, a group of morphologically heterogeneous neoplasms that share the same cytogenetic features. They are considered neural tumors derived from cells of the neural crest. Ewing sarcoma represents the less differentiated form of the tumors. Note=A chromosomal aberration involving ERG is found in patients with Erwing sarcoma. Translocation t(21;22)(q22;q12) with EWSR1. Note=Chromosomal aberrations involving ERG have been found in acute myeloid leukemia (AML). Translocation t(16;21)(p11;q22) with FUS. Translocation t(X;21)(q25-26;q22) with ELF4.
序列相似性	Belongs to the ETS family. Contains 1 ETS DNA-binding domain. Contains 1 PNT (pointed) domain.
细胞定位	Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Immunohistochemical analysis of formalin-fixed paraffin-embedded human kidney labelling ERG with ab92513 at a concentration of 1 µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with an OptiView DAB IHC Detection Kit. Heat mediated antigen retrieval was conducted for 32min with ULTRA cell conditioning solution (CC1 pH8.5). ab92513 anti ERG antibody was incubated at 37°C for 16min. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-ERG antibody [EPR3864] (ab92513)

Flow cytometry overlay histogram showing left THP-1 positive cells and right negative HCT116 stained with ab92513 (red line). The cells were fixed with 4% formaldehyde (10 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10 µg/ml human IgG and 10% normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab92513) (1x 10⁶ in 100 µl at 0.04 µg/ml (1/54000)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody (black line) was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.


Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.

Tissue Microarray (TMA) data for ab92513			
Normal tissue samples		Malignant tissue samples	
Human cardiac muscle	x (stromal cells ✓)	Human placenta	x (stromal cells ✓)
Human cerebrum	x (stromal cells ✓)	Human skeletal muscle	x (stromal cells ✓)
Human colon	x (stromal cells ✓)	Human skin	x (stromal cells ✓)
Human endometrium	x (stromal cells ✓)	Human spleen	✓
Human kidney	x (stromal cells ✓)	Human stomach	x (stromal cells ✓)
Human liver	x (stromal cells ✓)	Human testis	x (stromal cells ✓)
Human lung	✓	Human thyroid	x (stromal cells ✓)
Human mammary gland	x (stromal cells ✓)	Human tonsil	✓
Human pancreas	x (stromal cells ✓)		
		Clear cell carcinoma of human kidney	x (stromal cells ✓)
		Human bladder cancer	x (stromal cells ✓)
		Human breast carcinoma	x (stromal cells ✓)
		Human cervical carcinoma	x (stromal cells ✓)
		Human colon carcinoma	x (stromal cells ✓)
		Human endometrial carcinoma	x (stromal cells ✓)
		Human gastric adenocarcinoma	x (stromal cells ✓)
		Human prostatic adenocarcinoma	✓
		Human glioma	x (stromal cells ✓)
		Human hepatocellular carcinoma	x (stromal cells ✓)
		Human lung carcinoma	x (stromal cells ✓)
		Human ovarian carcinoma	x (stromal cells ✓)
		Human pancreatic carcinoma	x (stromal cells ✓)
		Human prostatic hyperplasia	x (stromal cells ✓)
		Human thyroid carcinoma	x (stromal cells ✓)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Tissue Microarrays stained for Anti-ERG antibody [EPR3864] using ab92513 in immunohistochemical analysis. This table provides a detailed overview of positive (tick mark) and negative (cross mark) staining per sample type tested. The section was incubated with ab92513 for 30 mins at room temperature followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). The immunostaining was performed on a Leica Biosystems BOND® RX instrument.


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 20 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Formalin-fixed, paraffin-embedded mouse brain tissue stained for ERG using ab92513 at 1/200 dilution in immunohistochemical analysis. A horse radish peroxidase antibody was used as the secondary antibody.

Antigen Retrieval: 40x; Proteinase K antigen retrieval - 15 min at 37 C



Western blot - Anti-ERG antibody [EPR3864]
(ab92513)

All lanes : purified

Lane 1 : rat brain lysate

Lane 2 : rat heart lysate

Lane 3 : RAW 264.7 (mouse macrophage cell line transformed with Abelson murine leukemia virus) cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 55 kDa

Observed band size: 55 kDa


Blocking buffer: 5% NFD/MTBST

Dilution buffer: 5% NFD/MTBST



Immunocytochemistry/ Immunofluorescence - Anti-
ERG antibody [EPR3864] (ab92513)


ab92513 staining ERG in THP-1 cells, with negative expression in HCT116 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised with 0.1% Triton x-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab92513 at 1 µg/ml and **ab7291**, Mouse monoclonal [DM1A] to alpha Tubulin at 0.5 µg/ml. Cells were then incubated with **ab150081**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and **ab150119**, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 647), pre-adsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue). Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.



Immunocytochemistry/ Immunofluorescence - Anti-
ERG antibody [EPR3864] (ab92513)

Image from KirbyBJ et al., PLoS One. 2012;8(12):e83903. Fig 4.; doi: 10.1371/journal.pone.0035976. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>

Functional characterization and detection of genetic alterations in GEDI-captured cells. The TMPRSS2:ERG fusion protein is detected in GEDI-captured circulating tumor cells (CTCs) from a castrate-resistant prostate cancer (CRPC) patient. PSMA-captured CTCs were stained on the device with ab92513. Representative examples of PSMA+/CD45- CTCs are shown, two of which are positive for ERG. Scale bars: 10 microns.




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-ERG antibody [EPR3864]
(ab92513)

Image from Litovkin K et al., PLoS One. 2015;10(6):e0130651. Fig 5.; doi: 10.1371/journal.pone.0130651. Reproduced under the Creative Commons license <https://creativecommons.org/publicdomain/zero/1.0/>


ERG and GSTP1 immunostainings of human prostate cancer samples using ab92513.

Representative immunohistochemical images of prostate cancer samples are shown that were positive for ERG and negative for GSTP1 (A), positive for both ERG and GSTP1 (B), negative for both ERG and GSTP1(C), and negative for ERG and positive for GSTP1 (D). The internal staining control for ERG is the endothelium (arrows) and for GSTP1 the stromal and/or basal cells of normal prostate glands. N, normal prostate gland; S, Stroma; T, tumor gland. Scale bars equal 100µm




Flow Cytometry (Intracellular) - Anti-ERG antibody [EPR3864] (ab92513)

Intracellular Flow Cytometry analysis of THP-1 (human monocytic leukemia cell line) cells labeling ERG with purified ab92513 at 1/1000 dilution (1ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) ([ab150077](#)) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) ([ab172730](#)) was used as the isotype control, Cell without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Immunohistochemical staining of paraffin embedded human kidney with purified ab92513 at a working dilution of 1/1000. The secondary antibody used is HRP goat anti-rabbit IgG H&L ([ab97051](#)) at 1/500. The sample is counter-stained with hematoxylin. Antigen retrieval was performed using Tris-EDTA buffer, pH 9.0. PBS was used instead of the primary antibody as the negative control, and is shown in the inset.



Western blot - Anti-ERG antibody [EPR3864] (ab92513)

All lanes : Anti-ERG antibody [EPR3864] (ab92513) at 1/2000 dilution (purified)

Lane 1 : Jurkat (human T cell leukemia cell line from peripheral blood) cell lysate

Lane 2 : HeLa (human epithelial cell line from cervix adenocarcinoma) cell lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : HRP goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 55 kDa

Observed band size: 55 kDa

Blocking buffer: 5% NFDm/TBST


Dilution buffer: 5% NFDm/TBST



Immunocytochemistry/ Immunofluorescence - Anti-ERG antibody [EPR3864] (ab92513)

Immunofluorescence staining of THP-1 (human monocytic leukemia cell line) cells with purified ab92513 at a working dilution of 1/100, counter-stained with DAPI. The secondary antibody was Alexa Fluor® 488 goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. **ab7291**, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with **ab150120** (Alexa Fluor® 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified ab92513 was used at a dilution of 1/500 followed by an Alexa Fluor® 594 goat anti-mouse antibody (**ab150120**) at a dilution of 1/500. For negative control 2, **ab7291** (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor® 488 goat anti-rabbit antibody (**ab150077**) at a dilution of 1/400.

Alexa Fluor[®] 488 (**ab196374**) and Alexa Fluor[®] 647 (**ab196149**) conjugated versions are available for this clone.




Western blot - Anti-ERG antibody [EPR3864] (ab92513)

Anti-ERG antibody [EPR3864] (ab92513) at 1/1000 dilution (unpurified) + Jurkat (human T cell leukemia cell line from peripheral blood) cell lysate at 10 µg

Secondary

HRP labelled Goat anti-Rabbit at 1/2000 dilution

Predicted band size: 55 kDa



Western blot - Anti-ERG antibody [EPR3864] (ab92513)

Lanes 1 & 3 : Anti-ERG antibody [EPR3864] (ab92513) at 1/250 dilution (unpurified)

Lanes 2 & 4 : Anti-ERG antibody [EPR3864] (ab92513) at 1/1000 dilution (unpurified)

Lane 1 : Jurkat (human T cell leukemia cell line from peripheral blood) Whole Cell Lysate

Lanes 2-4 : Jurkat Whole Cell Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/10000 dilution


Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 55 kDa

Observed band size: 55 kDa


Exposure time: 12 minutes



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ERG antibody [EPR3864] (ab92513)

Immunohistochemical analysis of paraffin embedded Human Prostatic adenocarcinoma stage 3 tissue using unpurified ab92513 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



OI-RD Scanning - Anti-ERG antibody [EPR3864] (ab92513)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ERG antibody [EPR3864] (ab92513)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors