

Anti-ERK5 antibody [EP791Y] ab40809

敲除验证
重组
RabMAb

★★★★★
[1 Abreviews](#)
[14 References](#)
[9 图像](#)

概述

产品名称	Anti-ERK5抗体[EP791Y]
描述	兔单克隆抗体[EP791Y] to ERK5
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IP, ICC/IF
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide within Human ERK5 aa 800-900 (C terminal). The exact sequence is proprietary.
阳性对照	WB: HeLa, HAP1, NIH/3T3 and PC-12 cell lysates. Flow Cyt (intra): HeLa and A549 cells. ICC: HeLa cells. IP: HeLa cell lysate.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS</p>
纯度	Protein A purified
克隆	单克隆
克隆编号	EP791Y

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab40809于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/50. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. For unpurified use at 1/100 dilution.
WB	★★★★★ (1)	1/10000. Detects a band of approximately 115 kDa (predicted molecular weight: 89 kDa). For unpurified use at 1/1000 - 1/5000 dilution.
IP		1/30. For unpurified use at 1/50 dilution.
ICC/IF		1/100. For unpurified use at 1/250 - 1/500 dilution.

靶标

功能	Plays a role in various cellular processes such as proliferation, differentiation and cell survival. The upstream activator of MAPK7 is the MAPK kinase MAP2K5. Upon activation, it translocates to the nucleus and phosphorylates various downstream targets including MEF2C. EGF activates MAPK7 through a Ras-independent and MAP2K5-dependent pathway. May have a role in muscle cell differentiation. May be important for endothelial function and maintenance of blood vessel integrity. MAP2K5 and MAPK7 interact specifically with one another and not with MEK1/ERK1 or MEK2/ERK2 pathways.
组织特异性	Expressed in many adult tissues. Abundant in heart, placenta, lung, kidney and skeletal muscle. Not detectable in liver.
序列相似性	Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily. Contains 1 protein kinase domain.
结构域	The second proline-rich region may interact with actin targeting the kinase to a specific location in the cell. The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.
翻译后修饰	Dually phosphorylated on Thr-219 and Tyr-221, which activates the enzyme (By similarity). Autophosphorylated in vitro on threonine and tyrosine residues when the C-terminal part of the kinase, which could have a regulatory role, is absent.
细胞定位	Cytoplasm. Nucleus. Translocates to the nucleus upon activation.

图片



Western blot - Anti-ERK5 antibody [EP791Y]
(ab40809)

All lanes : Anti-ERK5 antibody [EP791Y] (ab40809) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : MAPK7 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 89 kDa

Observed band size: 115 kDa


Lanes 1-2: Merged signal (red and green). Green - ab40809 observed at 115 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) observed at 37 kDa.

ab40809 was shown to react with ERK5 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line [ab265508](#) (knockout cell lysate [ab258042](#)) was used. Wild-type HeLa and MAPK7 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab40809 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-ERK5 antibody [EP791Y] (ab40809)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ERK5 with Purified ab40809 at 1:100 (4.8 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-ERK5 antibody [EP791Y] (ab40809)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling ERK5 with purified ab40809 at 1/50 dilution (10 µg/ml) (red). Cells were fixed with 80% methanol. A Goat anti rabbit IgG (Alexa Fluor®488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-ERK5 antibody [EP791Y] (ab40809)

All lanes : Anti-ERK5 antibody [EP791Y] (ab40809) at 1/2000 dilution (Purified)

Lane 1 : HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates with 5% NFD/MTBST

Lane 2 : NIH/3T3 (Mouse embryonic fibroblast) whole cell lysates with 5% NFD/MTBST

Lane 3 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates with 5% NFD/MTBST


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 89 kDa

Observed band size: 115 kDa



Western blot - Anti-ERK5 antibody [EP791Y]
(ab40809)


Lane 1: Wild-type HAP1 whole cell lysate (20 μ g)

Lane 2: MAPK7 (ERK5) knockout HAP1 whole cell lysate (20 μ g)

Lane 3: HeLa whole cell lysate (20 μ g)

Lanes 1 - 3: Merged signal (red and green). Green - ab40809 observed at 88 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

Unpurified ab40809 was shown to specifically react with ERK5 in wild-type HAP1 cells as signal was lost in MAPK7 (ERK5) knockout cells. Wild-type and MAPK7 (ERK5) knockout samples were subjected to SDS-PAGE. Unpurified ab40809 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Western blot - Anti-ERK5 antibody [EP791Y]
(ab40809)

Anti-ERK5 antibody [EP791Y] (ab40809) at 1/5000 dilution (unpurified) + HeLa cell lysate at 10 μ g

Predicted band size: 89 kDa


Observed band size: 115 kDa

The predicted weight of 89 kDa is for the precursor version of human ERK5 protein. However, ab40809 detects endogenous levels of total Erk5 protein which appears around 115 kDa in SDS PAGE



Flow Cytometry (Intracellular) - Anti-ERK5 antibody [EP791Y] (ab40809)

Overlay histogram showing A549 cells stained with unpurified ab40809 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab40809, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor[®] 488 goat anti-rabbit IgG (H&L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunoprecipitation - Anti-ERK5 antibody [EP791Y] (ab40809)

ab40809 (purified) at 1:30 dilution (2ug) immunoprecipitating ERK5 in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10ug

Lane 2 (+): ab40809 & HeLa whole cell lysate

Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab40809 in HeLa whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ERK5 antibody [EP791Y] (ab40809)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors