

Anti-Factor VIII antibody [GMA-012] ab78852

4 References 2 图像

概述	
产品名称	Anti-Factor VIII抗体[GMA-012]
描述	小鼠单克隆抗体[GMA-012] to Factor VIII
宿主	Mouse
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human
免疫原	Purified human Factor VIII
阳性对照	Human Factor VIII. This antibody gave a positive result in IHC in the following FFPE tissue: Human normal liver.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
性能	
形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 7.40 Constituents: 1% Mannitol, 0.87% Sodium chloride, 0.164% Sodium phosphate
纯度	DEAE-Chromatography
克隆	单克隆
克隆编号	GMA-012
同种型	IgG1
应用	


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use a concentration of 1 µg/ml. Detects a band of approximately 280 kDa (predicted molecular weight: 267 kDa).
IHC-P		Use a concentration of 5 µg/ml.

靶标

功能	Factor VIII, along with calcium and phospholipid, acts as a cofactor for factor IXa when it converts factor X to the activated form, factor Xa.
疾病相关	Defects in F8 are the cause of hemophilia A (HEMA) [MIM:306700]. A disorder of blood coagulation characterized by a permanent tendency to hemorrhage. About 50% of patients have severe hemophilia resulting in frequent spontaneous bleeding into joints, muscles and internal organs. Less severe forms are characterized by bleeding after trauma or surgery. Note=Of particular interest for the understanding of the function of F8 is the category of CRM (cross-reacting material) positive patients (approximately 5%) that have considerable amount of F8 in their plasma (at least 30% of normal), but the protein is non-functional; i.e., the F8 activity is much less than the plasma protein level. CRM-reduced is another category of patients in which the F8C antigen and activity are reduced to approximately the same level. Most mutations are CRM negative, and probably affect the folding and stability of the protein.
序列相似性	Belongs to the multicopper oxidase family. Contains 3 F5/8 type A domains. Contains 2 F5/8 type C domains. Contains 6 plastocyanin-like domains.
结构域	Domain F5/8 type C 2 is responsible for phospholipid-binding and essential for factor VIII activity.
翻译后修饰	Sulfation on Tyr-1699 is essential for binding vWF.
细胞定位	Secreted > extracellular space.

图片




Anti-Factor VIII antibody [GMA-012] (ab78852) at 1 µg/ml + human Factor VIII at 2 µg

Secondary

Goat anti-mouse HRP conjugated IgG

Predicted band size: 267 kDa

Observed band size: 280 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Factor VIII antibody [GMA-012] (ab78852)

IHC image of Factor VIII staining in Human normal liver formalin fixed paraffin embedded tissue section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab78852, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors