

### Anti-FATS antibody ab122497

[1 References](#) [3 图像](#)

#### 概述

产品名称	Anti-FATS抗体
描述	兔多克隆抗体to FATS
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Human
免疫原	Recombinant fragment corresponding to Human FATS aa 169-262. Sequence: AEDTLFQAPPALANGAHPGRHQRSFACTEF SRNSSVRLKVP EAHTGLCE RRKYVWTHADDKETSFSPDTPLSGKSPLVFSSCVHLRVSQQC PD <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
阳性对照	Human kidney tissue; RT-4 and U-251 MG cell lysate; Human Liver and Tonsil tissue lysate.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

## 应用


**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab122497于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use a concentration of 1 - 4 µg/ml. Recommend PFA Fixation and Triton X-100 treatment
IHC-P		1/200 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		1/250 - 1/500.


## 靶标

## 图片




ab122497 at 1/425 staining FATS in Paraffin Embedded Human kidney tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FATS antibody (ab122497)



Immunocytochemistry/ Immunofluorescence - Anti-FATS antibody (ab122497)

Immunofluorescent staining of Human cell line A-431 shows positivity in nucleus but not nucleoli, plasma membrane and cytoplasm. Recommended concentration of ab122497 1-4 µg/ml. Cells treated with PFA/Triton X-100.



Western blot - Anti-FATS antibody (ab122497)

**All lanes** : Anti-FATS antibody (ab122497) at 1/250 dilution

**Lane 1** : RT-4 lysate

**Lane 2** : U-251 MG lysate

**Lane 3** : Human Plasma

**Lane 4** : Human Liver lysate

**Lane 5** : Human Tonsil lysate

Developed using the ECL technique.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors