

# Anti-FGF17 antibody ab9990

## 1 图像

### 概述

---

产品名称	Anti-FGF17抗体
描述	兔多克隆抗体to FGF17
宿主	Rabbit
经测试应用	适用于: Sandwich ELISA
免疫原	Recombinant full length protein corresponding to Human FGF17 aa 1-250. Database link: <a href="#">O60258</a> <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
阳性对照	Recombinant human FGF17 protein ( <a href="#">ab9991</a> ) can be used as a positive control in WB.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

---

形式	Lyophilized:Reconstitute with 200µl of sterile water. Please note that if you receive this product in liquid form it has already been reconstituted as described and no further reconstitution is necessary.
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	No preservative, sterile filtered
纯度	Immunogen affinity purified
克隆	多克隆
同种型	unknown
轻链类型	unknown

### 应用

---

## The Abpromise guarantee

**Abpromise™** 承诺保证使用 ab9990 于以下的经测试应用


“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Sandwich ELISA		Use at an assay dependent concentration. Can be paired for Sandwich ELISA with <b>Mouse monoclonal [MM0284-5H27] to FGF17 (ab89270)</b> . For sandwich ELISA, use this antibody as Detection at 0.5µg/ml with <b>ab89270</b> as Capture.

## 靶标

功能	Plays an important role in the regulation of embryonic development and as signaling molecule in the induction and patterning of the embryonic brain. Required for normal brain development.
组织特异性	Preferentially expressed in the embryonic brain.
序列相似性	Belongs to the heparin-binding growth factors family.
发展阶段	Detected in embryos at E14.5, but not at E10.5 and E19.5. Preferentially expressed in the neuroepithelia of the isthmus and septum of the embryonic brain at E14.5.
细胞定位	Secreted.

## 图片



Standard Curve for FGF17 (Analyte: **ab9991**) dilution range 1pg/ml to 1ug/ml using Capture Antibody Mouse monoclonal [MM0284-5H27] to FGF17 (**ab89270**) at 1ug/ml and Detector Antibody Rabbit polyclonal to FGF17 (ab9990) at 0.5ug/ml

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors