

Anti-FHL2 antibody [EPR17860-20] ab202584

敲除验证 重组 RabMAb

4 References 11 图像

概述

| | |
|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 产品名称 | Anti-FHL2抗体[EPR17860-20] |
| 描述 | 兔单克隆抗体[EPR17860-20] to FHL2 |
| 宿主 | Rabbit |
| 经测试应用 | 适用于: WB, ICC/IF, IP |
| 种属反应性 | 与反应: Mouse, Rat, Human |
| 免疫原 | Recombinant fragment. This information is proprietary to Abcam and/or its suppliers. |
| 阳性对照 | WB: U-2 OS, K526, HeLa, SW480, PC-12 and HT1080 whole cell lysates; Human fetal heart lysate; Mouse and rat heart lysates. ICC/IF: A-673 and NIH/3T3 cells. IP: SW480 whole cell lysate. ICC/IF: U-2 OS cells. |
| 常规说明 | <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> |

性能

| | |
|------|-----------------------------------------------------------------------------------------------------------------------------------|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. |
| 存储溶液 | pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol, 0.05% BSA |
| 纯度 | Protein A purified |
| 克隆 | 单克隆 |
| 克隆编号 | EPR17860-20 |

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab202584于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|--------|------|---------------------------------------------------------------------------------------------------------------------|
| WB | | 1/1000. Detects a band of approximately 32 kDa (predicted molecular weight: 32 kDa). |
| ICC/IF | | 1/400 - 1/200. ab202584 works both with PFA and methanol fixation. Fixation with PFA gives the strongest signal. |
| IP | | 1/30. |

靶标

功能

May function as a molecular transmitter linking various signaling pathways to transcriptional regulation. Negatively regulates the transcriptional repressor E4F1 and may function in cell growth.

组织特异性

Expressed in skeletal muscle and heart.

序列相似性

Contains 4 LIM zinc-binding domains.


结构域

The third LIM zinc-binding mediates interaction with E4F1.

细胞定位

Cytoplasm. Nucleus.

图片



Western blot - Anti-FHL2 antibody [EPR17860-20]
(ab202584)

All lanes : Anti-FHL2 antibody [EPR17860-20] (ab202584) at
1/1000 dilution

Lane 1 : Wild-type U-2 OS cell lysate

Lane 2 : FHL2 knockout U-2 OS cell lysate

Lane 3 : HeLa (Human epithelial cell line from cervix
adenocarcinoma) whole cell lysate

Lysates/proteins at 40 µg per lane.


Performed under reducing conditions.

Predicted band size: 32 kDa

Observed band size: 32 kDa

Lanes 1 - 3: Merged signal (red and green). Green - ab202584 observed at 32 kDa. Red - loading control **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) observed at 55kDa.


ab202584 was shown to react with FHL2 in wild-type U-2 OS cells in western blot with loss of signal observed in FHL2 knockout sample. Wild-type and FHL2 knockout U-2 OS cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween[®]) before incubation with ab202584 and **ab7291** (Mouse anti-Alpha Tubulin [DM1A]) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye[®] 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye[®] 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-FHL2 antibody [EPR17860-20] (ab202584)

ab202584 staining FHL2 in wild-type U2OS cells (top panel) and FHL2 knockout U2OS cells (bottom panel). The cells were fixed with 4% PFA (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab202584 at 1/200 dilution and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor[®] 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor[®] 594) (**ab150120**) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-FHL2 antibody [EPR17860-20] (ab202584)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized A-673 (Human muscle Ewing's Sarcoma cell line) cells labeling FHL2 with ab202584 at 1/400 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green). Confocal image showing cytoplasmic and weakly nuclear staining on A-673 cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: ab202584 at 1/400 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.



Western blot - Anti-FHL2 antibody [EPR17860-20] (ab202584)

All lanes : Anti-FHL2 antibody [EPR17860-20] (ab202584) at 1/1000 dilution

Lane 1 : Wild-type HeLa lysate

Lane 2 : FHL2 knockout HeLa lysate

Lane 3 : K562 lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 32 kDa

Lanes 1-3: Merged signal (red and green). Green - ab202584 observed at 32 kDa. Red - loading control **ab7291** observed at 50 kDa.

ab202584 Recombinant Anti-FHL2 antibody [EPR17860-20] was shown to specifically react with FHL2 in wild-type HeLa cells. Loss of signal was observed when knockout cell line **ab265475** (knockout cell lysate **ab257441**) was used. Wild-type and FHL2 knockout samples were subjected to SDS-PAGE. ab202584 and Anti-alpha Tubulin antibody [DM1A] - Loading Control? (**ab7291**) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and


Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry/ Immunofluorescence - Anti-FHL2 antibody [EPR17860-20] (ab202584)

ab202584 staining FHL2 in wild-type U2OS cells (top panel) and FHL2 knockout U2OS cells (bottom panel). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab202584 at 1/200 dilution and **ab7291** (Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-FHL2 antibody [EPR17860-20] (ab202584)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryo fibroblast cells) cells labeling FHL2 with ab202584 at 1/400 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/500 dilution (green). Confocal image showing cytoplasmic and weakly nuclear staining on NIH/3T3 cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution and **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).

The negative controls are as follows:

-ve control 1: ab202584 at 1/400 dilution followed by **ab150120** (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.

-ve control 2: **ab7291** (anti-Tubulin mouse mAb) at 1/1000 dilution followed by **ab150077** (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.



Western blot - Anti-FHL2 antibody [EPR17860-20]
(ab202584)

All lanes : Anti-FHL2 antibody [EPR17860-20] (ab202584) at 1/1000 dilution

Lane 1 : SW480 (Human colorectal adenocarcinoma cell line) whole cell lysate

Lane 2 : PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate

Lane 3 : HT1080 (Human fibrosarcoma cells) whole cell lysate

Lane 4 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/1000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 32 kDa

Observed band size: 32 kDa

Exposure time: 30 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-FHL2 antibody [EPR17860-20] (ab202584)

Anti-FHL2 antibody [EPR17860-20] (ab202584) at 1/10000 dilution
+ Human fetal heart lysate at 10 µg

Secondary


Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 32 kDa

Observed band size: 32 kDa

Exposure time: 15 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot - Anti-FHL2 antibody [EPR17860-20] (ab202584)

All lanes : Anti-FHL2 antibody [EPR17860-20] (ab202584) at 1/2000 dilution

Lane 1 : Mouse heart lysate

Lane 2 : Rat heart lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/1000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 32 kDa

Observed band size: 32 kDa

Exposure time: 10 seconds

Blocking/Dilution buffer: 5% NFDm/TBST.



Immunoprecipitation - Anti-FHL2 antibody
[EPR17860-20] (ab202584)



FHL2 was immunoprecipitated from 1mg of SW480 (Human colorectal adenocarcinoma cell line) whole cell lysate with ab202584 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab202584 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution.

Lane 1: SW480 whole cell lysate 10 µg (Input). Lane 2: ab202584 IP in SW480 whole cell lysate. Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of ab202584 in SW480 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDm/TBST.

Exposure time: 10 seconds.

Why choose a recombinant antibody?

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|
|  Research with confidence Consistent and reproducible results |  Long-term and scalable supply Recombinant technology |
|  Success from the first experiment Confirmed specificity |  Ethical standards compliant Animal-free production |

Anti-FHL2 antibody [EPR17860-20] (ab202584)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors