

FITC Anti-CD146 antibody [OJ79c] ab33300

2 References

概述

产品名称	FITC荧光Anti-CD146抗体[OJ79c]
描述	FITC荧光小鼠单克隆抗体[OJ79c] to CD146
宿主	Mouse
偶联物	FITC. Ex: 493nm, Em: 528nm
特异性	Recognises the 118kD cell surface glycoprotein CD146, also known as MUC18, MelCAM and Sendo
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Human
免疫原	Recombinant MUC18 (D1-D5) Fc protein (Human)
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. Store In the Dark.
存储溶液	Preservative: 0.09% Sodium azide Constituents: PBS, 1% BSA
纯度	Protein G purified
纯化说明	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
克隆	单克隆
克隆编号	OJ79c
骨髓瘤	Sp2/0-Ag14
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab33300于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use at an assay dependent concentration. or 100ul whole blood. ab91356 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

靶标

功能	Plays a role in cell adhesion, and in cohesion of the endothelial monolayer at intercellular junctions in vascular tissue. Its expression may allow melanoma cells to interact with cellular elements of the vascular system, thereby enhancing hematogeneous tumor spread. Could be an adhesion molecule active in neural crest cells during embryonic development. Acts as surface receptor that triggers tyrosine phosphorylation of FYN and PTK2, and a transient increase in the intracellular calcium concentration.
组织特异性	Detected in endothelial cells in vascular tissue throughout the body. May appear at the surface of neural crest cells during their embryonic migration. Appears to be limited to vascular smooth muscle in normal adult tissues. Associated with tumor progression and the development of metastasis in human malignant melanoma. Expressed most strongly on metastatic lesions and advanced primary tumors and is only rarely detected in benign melanocytic nevi and thin primary melanomas with a low probability of metastasis.
序列相似性	Contains 3 Ig-like C2-type (immunoglobulin-like) domains. Contains 2 Ig-like V-type (immunoglobulin-like) domains.
细胞定位	Membrane.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors