


# FITC Anti-CD177 antibody [MEM-166] ab26013

**3 References**   **1 图像**

### 概述

|       |   |
|-------|---|
| 产品名称  | FITC荧光Anti-CD177抗体[MEM-166]   |
| 描述    | FITC荧光小鼠单克隆抗体[MEM-166] to CD177   |
| 宿主    | Mouse   |
| 偶联物   | FITC. Ex: 493nm, Em: 528nm  |
| 经测试应用 | 适用于: Flow Cyt   |
| 种属反应性 | 与反应: Human<br>预测可用于: Non human primates    |
| 免疫原   | Tissue, cells or virus corresponding to Human CD177. Human granulocytes   |
| 常规说明  | <p>The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC and adjusted for direct use.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

### 性能

|      |  |
|------|--|
| 形式   | Liquid   |
| 存放说明 | Shipped at 4°C. Store at +4°C.   |
| 存储溶液 | <p>pH: 7.4</p> <p>Preservative: 0.097% Sodium azide</p> <p>Constituents: PBS, 0.2% BSA</p> |
| 纯度   | Size exclusion   |
| 纯化说明 | Purity >95% by SDS-PAGE.   |
| 克隆   | 单克隆  |
| 克隆编号 | MEM-166  |
| 同种型  | IgG1   |

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab26013于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用       | Ab评论 | 说明  |
|----------|------|---|
| Flow Cyt |      | Use 20µl for 10 <sup>6</sup> cells.<br>or 100 µl of whole blood. (The antibody should be used undiluted with the samples).<br><br><b>ab91356</b> - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody. |

靶标

|       |  |
|-------|--|
| 组织特异性 | Highly expressed in normal bone marrow and weakly expressed in fetal liver. Expressed on neutrophils. Expressed in granulocytes of patients with polycythemia vera (PV) and with essential thrombocythemia (ET). |
| 序列相似性 | Contains 2 UPAR/Ly6 domains.   |
| 翻译后修饰 | N-glycosylated.<br>A soluble form may also be produced by proteolytic cleavage at the cell surface (shedding).   |
| 细胞定位  | Cell membrane.   |

图片



Surface staining of human peripheral blood cells with anti-CD177.  
ab26013 was used at a recommended concentration of 1 µg/mL.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors