

FITC Anti-CD52 antibody [HI186], prediluted ab69774

2 图像

概述

| | |
|--------------|---|
| 产品名称 | FITC荧光Anti-CD52抗体[HI186], prediluted |
| 描述 | FITC荧光小鼠单克隆抗体[HI186] to CD52, prediluted |
| 宿主 | Mouse |
| 偶联物 | FITC. Ex: 493nm, Em: 528nm |
| 经测试应用 | 适用于: Flow Cyt |
| 种属反应性 | 与反应: Human |
| 免疫原 | Tissue, cells or virus corresponding to Human CD52. Human tonsil |
| 常规说明 | <p>The purified antibody is conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The reagent is free of unconjugated FITC.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

性能

| | |
|-------------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C. |
| 存储溶液 | pH: 7.4 Preservative: 0.097% Sodium azide Constituents: 0.2% BSA, PBS |
| 纯度 | Size exclusion |
| 克隆 | 单克隆 |
| 克隆编号 | HI186 |
| 同种型 | IgG2b |

The Abpromise guarantee **Abpromise™**承诺保证使用ab69774于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|----------|------|--|
| Flow Cyt | | Use 5µl for 10 ⁶ cells. ab91368 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody. |

靶标


| | |
|-------------|---|
| 功能 | May play a role in carrying and orienting carbohydrate, as well as having a more specific role. |
| 细胞定位 | Cell membrane. |

图片



Overlay histogram showing peripheral blood lymphocytes stained with ab69774 (red line). The cells incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab69774, 0.06 µg/1x10⁶ cells) for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2b FITC (0.6 µg/1x10⁶ cells) for 30 min at 22°C. Acquisition of >5,000 events was performed.

Flow Cytometry - FITC Anti-CD52 antibody [HI186], prediluted (ab69774)



Surface staining of CD52 in human peripheral blood with ab69774

Flow Cytometry - FITC Anti-CD52 antibody [HI186], prediluted (ab69774)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors