

FITC Anti-Dectin-1 antibody [2A11] ab21646

★★★★★ [1 Abreviews](#) [1 References](#) [1 图像](#)

概述

产品名称	FITC荧光Anti-Dectin-1抗体[2A11]
描述	FITC荧光大鼠单克隆抗体[2A11] to Dectin-1
宿主	Rat
偶联物	FITC. Ex: 493nm, Em: 528nm
经测试应用	适用于: Flow Cyt
种属反应性	与反应: Mouse
免疫原	Recombinant full length protein. This information is proprietary to Abcam and/or its suppliers.
常规说明	<p>This antibody inhibits the binding of zymosan to macrophages via the beta Glucan Receptor.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.09% Sodium azide</p> <p>Constituents: PBS, 1% BSA</p>
纯度	Protein G purified
Primary antibody说明	This antibody inhibits the binding of zymosan to macrophages via the beta Glucan Receptor.
克隆	单克隆
克隆编号	2A11
骨髓瘤	Y3/Ag1.2.3

同种型

IgG2b

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab21646于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt	★★★★★ (1)	Use 10µl for 10 ⁶ cells. Use 10µl for 106 cells in 100µl.

靶标

功能	Lectin that functions as pattern receptor specific for beta-1,3-linked and beta-1,6-linked glucans, such as cell wall constituents from pathogenic bacteria and fungi. Necessary for the TLR2-mediated inflammatory response and for TLR2-mediated activation of NF-kappa-B. Enhances cytokine production in macrophages and dendritic cells. Mediates production of reactive oxygen species in the cell. Mediates phagocytosis of C.albicans conidia. Binds T-cells in a way that does not involve their surface glycans and plays a role in T-cell activation. Stimulates T-cell proliferation.
组织特异性	Highly expressed in peripheral blood leukocytes and dendritic cells. Detected in spleen, bone marrow, lung, muscle, stomach and placenta.
疾病相关	Defects in CLEC7A may be a cause of familial candidiasis type 4 (CANDF4) [MIM:613108]. It is a rare disorder with altered immune responses and impaired clearance of fungal infections, selective against Candida. It is characterized by persistent and/or recurrent infections of the skin, nails and mucous membranes caused by organisms of the genus Candida, mainly Candida albicans.
序列相似性	Contains 1 C-type lectin domain.
翻译后修饰	Phosphorylated on tyrosine residues in response to glucan binding.
细胞定位	Cytoplasm and Cell membrane.

图片



Flow cytometry staining of mouse peripheral blood granulocytes with ab21646.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors