


Anti-FOXA2 antibody [EPR4466] ab108422

重组 RabMAb

★★★★★ **10 Abreviews** **86 References** **10 图像**

概述

产品名称	Anti-FOXA2抗体[EPR4466]
描述	兔单克隆抗体[EPR4466] to FOXA2
宿主	Rabbit
经测试应用	适用于: WB, IHC-P, ICC/IF
种属反应性	与反应: Mouse, Human 预测可用于: Rat 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human colon cancer, fetal colon and mouse lung tissue lysates and HepG2 whole cell lysate (ab7900). IHC-P: Human hepatocellular carcinoma and mouse liver tissue; Human colon tissue; Mouse brain (substantia nigra) tissue. ICC/IF: HT-29 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol, 0.05% BSA, 59% PBS
纯度	Protein A purified
克隆	单克隆

同种型 IgG

The Abpromise guarantee

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Predicted molecular weight: 52 kDa. For unpurified use at 1/1000 - 1/10000.
IHC-P	★★★★★ (6)	1/500 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/250 - 1/500.
ICC/IF		1/300. For unpurified use at 1/250 - 1/500.


功能

Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). In embryonic development is required for notochord formation. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas and lungs; FOXA1 and FOXA2 seem to have at least in part redundant roles. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; regulates the expression of genes important for glucose sensing in pancreatic beta-cells and glucose homeostasis. Involved in regulation of fat metabolism. Binds to fibrinogen beta promoter and is involved in IL6-induced fibrinogen beta transcriptional activation.

Contains 1 fork-head DNA-binding domain.

Phosphorylation on Thr-156 abolishes binding to target promoters and subsequent transcription activation upon insulin stimulation.

Nucleus. Cytoplasm. Shuttles between the nucleus and cytoplasm in a CRM1-dependent manner and in response to insulin signaling via AKT1 is exported from the nucleus.




Immunocytochemistry/ Immunofluorescence - Anti-FOXA2 antibody [EPR4466] (ab108422)

Immunocytochemistry/Immunofluorescence analysis of HT-29 cells labelling FOXA2 with purified ab108422 at 1/300. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/500) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.


-ve control 1: primary antibody (1/300) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

-ve control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXA2 antibody [EPR4466] (ab108422)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human hepatocellular carcinoma tissue labelling FOXA2 with purified ab108422 at 1/500. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Western blot - Anti-FOXA2 antibody [EPR4466] (ab108422)

All lanes : Anti-FOXA2 antibody [EPR4466] (ab108422) at 1/1000 dilution (unpurified)

Lane 1 : Human fetal colon tissue lysate


Lane 2 : HepG2 (human liver hepatocellular carcinoma cell line) cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP-conjugated goat anti-rabbit IgG

Predicted band size: 52 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXA2 antibody [EPR4466] (ab108422)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of mouse liver tissue labelling FOXA2 with purified ab108422 at 1/500. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. [ab97051](#), a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.



Western blot - Anti-FOXA2 antibody [EPR4466]
(ab108422)

Anti-FOXA2 antibody [EPR4466] (ab108422) at 1/10000 dilution
(purified) + Human colon cancer tissue lysate at 20 µg

Secondary


Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 52 kDa

Observed band size: 52 kDa

Blocking buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-FOXA2 antibody
[EPR4466] (ab108422)

This image is courtesy of Carl Hobbs, King's College
London, United Kingdom

Immunohistochemical analysis of Formalin/PFA-fixed paraffin-
embedded human colon sections labelling FOXA2 with ab108422
at dilution of 1/500. The secondary antibody used was a polyclonal
goat anti-rabbit biotin conjugated antibody at a dilution of 1/300.
The sample was counterstained with hematoxylin. Antigen retrieval
was heat mediated using citric acid.



Western blot - Anti-FOXA2 antibody [EPR4466] (ab108422)

Anti-FOXA2 antibody [EPR4466] (ab108422) at 1/2000 dilution (purified) + Mouse lung tissue lysate at 20 µg

Secondary


Peroxidase-conjugated goat anti-rabbit IgG (H+L) at 1/1000 dilution

Predicted band size: 52 kDa

Observed band size: 52 kDa

Blocking buffer and concentration: 5% NFDM/TBST.


Diluting buffer and concentration: 5% NFDM /TBST.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXA2 antibody [EPR4466] (ab108422)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human hepatocellular carcinoma tissue labelling FOXA2 with unpurified ab108422 at a 1/250 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-FOXA2 antibody [EPR4466] (ab108422)

This image is courtesy of Carl Hobbs, King's College London, United Kingdom

ab108422 staining of FOXA2 in mouse brain (substantia nigra) tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Heat-mediated antigen retrieval was carried out using citric acid. Samples were incubated with primary antibody (1/500) for two hours at room temperature. A Biotin-conjugated goat anti-rabbit IgG polyclonal was used as the secondary antibody.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-FOXA2 antibody [EPR4466] (ab108422)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors