




### Anti-FOXP2 antibody - C-terminal ab1307

★★★★★ [7 Abreviews](#) [28 References](#) [1 图像](#)

#### 概述

|       |   |
|-------|---|
| 产品名称  | Anti-FOXP2抗体- C-terminal  |
| 描述    | 山羊多克隆抗体to FOXP2 - C-terminal  |
| 宿主    | Goat  |
| 特异性   | This antibody is expected to recognise all three reported isoforms (NP_055306.1; NP_683696.2; NP_683697.1).   |
| 经测试应用 | <b>适用于:</b> WB  |
| 种属反应性 | <b>与反应:</b> Human<br><b>预测可用于:</b> Mouse, Rat, Cow, Dog, Pig, Xenopus laevis, Chimpanzee, Rhesus monkey, Gorilla<br>   |
| 免疫原   | Synthetic peptide corresponding to Human FOXP2 aa 703-715 (C terminal) (Cysteine residue).<br>Sequence:<br>C-REIEEEPLSEDLE<br><br>Database link: <a href="#">O15409</a><br>(Peptide available as <a href="#">ab22800</a> )<br><div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div>  |
| 阳性对照  | WB: Human cerebellum cell lysate.   |
| 常规说明  | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p> |

#### 性能

|      |   |
|------|---|
| 形式   | Liquid  |
| 存放说明 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| 存储溶液 | pH: 7.30<br>Preservative: 0.02% Sodium azide  |

|      |   |
|------|---|
| 纯度   | Constituents: 0.05% Tris, 0.5% BSA  |
| 纯化说明 | Immunogen affinity purified   |
| 克隆   | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| 同种型  | 多克隆   |
|      | IgG   |

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab1307于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论      | 说明  |
|----|-----------|---|
| WB | ★★★★★ (1) | Use a concentration of 0.5 - 2 µg/ml. Detects a band of approximately 80 kDa (predicted molecular weight: 79.9 kDa). 1 hour primary incubation is recommended for this product. |

靶标

|       |   |
|-------|---|
| 功能    | Transcriptional repressor that may play a role in the specification and differentiation of lung epithelium. May also play a role in developing neural, gastrointestinal and cardiovascular tissues. Can act with CTBP1 to synergistically repress transcription but CTPBP1 is not essential. Involved in neural mechanisms mediating the development of speech and language.  |
| 组织特异性 | Isoform 1 and isoform 6 are expressed in adult and fetal brain, caudate nucleus and lung.   |
| 疾病相关  | Defects in FOXP2 are the cause of speech-language disorder 1 (SPCH1) [MIM:602081]; also known as autosomal dominant speech and language disorder with orofacial dyspraxia. Affected individuals have a severe impairment in the selection and sequencing of fine orofacial movements, which are necessary for articulation. They also show deficits in several facets of language processing (such as the ability to break up words into their constituent phonemes) and grammatical skills.<br>Note=A chromosomal aberration involving FOXP2 is a cause of severe speech and language impairment. Translocation t(5;7)(q22;q31.2).   |
| 序列相似性 | Contains 1 C2H2-type zinc finger.<br>Contains 1 fork-head DNA-binding domain.   |
| 发展阶段  | Expressed in the brain at 15 and 22 weeks of gestation, with a pattern of strong cortical, basal ganglia, thalamic and cerebellar expression. Highly expressed in the head and tail of nucleus caudatus and putamen. Restricted expression within the globus pallidus, with high levels in the pars interna, which provides the principal source of output from the basal ganglia to the nucleus centrum medianum thalami (CM) and the major motor relay nuclei of the thalamus. In the thalamus, present in the CM and nucleus medialis dorsalis thalami. Lower levels are observed in the nuclei anterior thalami, dorsal and ventral, and the nucleus parafascicularis thalami. Expressed in the ventrobasal complex comprising the nucleus ventralis posterior lateralis/medialis. The ventral tier of the thalamus exhibits strong expression, including nuclei ventralis anterior, lateralis and posterior lateralis pars oralis. Also expressed in the nucleus subthalamicus bilaterally and in the nucleus ruber. |


## 结构域

The leucine-zipper is required for dimerization and transcriptional repression.

## 细胞定位

Nucleus.

## 图片



Anti-FOXP2 antibody - C-terminal (ab1307) at 2 µg/ml + Human cerebellum whole cell lysate

Developed using the ECL technique.

**Predicted band size: 79.9 kDa**

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors