abcam

Product datasheet

Anti-GABARAP antibody ab1398

**** 1 Abreviews 4 References

概述

产品名称 Anti-GABARAP抗体

描述 兔多克隆抗体to GABARAP

宿主 Rabbit

特异性 This antibody recognizes human GABARAP (14 kD).

经测试应用 适用于: ELISA, WB

种属反应性 与反应: Mouse, Human

免疫原 Synthetic peptide:

RSEGEKIRKKYPDRVPV

, corresponding to amino acids 15-31 of Human GABARAP.

Run BLAST with

Run BLAST with

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

存储溶液 pH: 7.4

Preservative: 0.02% Sodium azide

纯**度** Affinity purified

应用

1

The Abpromise guarantee

Abpromise™承诺保证使用ab1398于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
WB	*** <u>*</u>	Use at an assay dependent concentration.

靶标

功能	Ubiquitin-like modifier that plays a role in intracellular transport of GABA(A) receptors and its interaction with the cytoskeleton. Involved in apoptosis. Involved in autophagy. Whereas LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16 subfamily is essential for a later stage in autophagosome maturation.	
组织 特异性	Heart, brain, placenta, liver, skeletal muscle, kidney and pancreas.	
序列相似性	Belongs to the ATG8 family.	
翻译后修饰	The precursor molecule is cleaved by ATG4B to form the cytosolic form, GABARAP-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, GABARAP-II.	
细胞定位	Endomembrane system. Cytoplasm, cytoskeleton. Golgi apparatus membrane. Cytoplasmic vesicle, autophagosome. Largely associated with intracellular membrane structures including the Golgi apparatus and postsynaptic cisternae. Colocalizes with microtubules (By similarity). Localizes also to discrete punctae along the ciliary axoneme (By similarity).	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors