

Anti-GAPDH antibody [EPR6256] - Loading Control ab128915

重组 RabMAb

★★★★★ **11 Abreviews** **246 References** **13 图像**

概述

产品名称	Anti-GAPDH抗体[EPR6256] - Loading Control
描述	兔单克隆抗体[EPR6256] to GAPDH - Loading Control
宿主	Rabbit
特异性	The African green monkey recommendation is based on the WB results. We do not guarantee IHC-P for African green monkey.
经测试应用	适用于: WB, IP, ICC/IF, IHC-P, Flow Cyt (Intra)
种属反应性	与反应: Human, African green monkey
免疫原	Synthetic peptide within Human GAPDH aa 250 to the C-terminus. The exact sequence is proprietary. Database link: P04406
阳性对照	WB: 293T, HeLa, HepG2, HUVEC, MCF7 and SH-SY5Y cell lysates. ICC/IF: HeLa and MCF7 cells. IHC-P: Human pancreas tissue, Human bladder carcinoma. Flow Cyt (intra): HeLa cells. IP: HeLa cell lysate.
常规说明	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents . Mouse: We have preliminary internal testing data to indicate this antibody may not react with this species. Please contact us for more information.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C.
存储溶液	pH: 7.20

	Preservative: 0.01% Sodium azide
	Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR6256
同种型	IgG

应用

The Abpromise guarantee
Abpromise™承诺保证使用ab128915于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (9)	1/10000 - 1/50000. Detects a band of approximately 35 kDa (predicted molecular weight: 36 kDa).
IP		1/10 - 1/100.
ICC/IF	★★★★★ (1)	1/250 - 1/500. 2.0 µg/ml
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. See IHC antigen retrieval protocols . For unpurified use at 1/250. The African green monkey recommendation is based on the WB results. We do not guarantee IHC-P for African green monkey.
Flow Cyt (Intra)		1/20. For unpurified use at 1/100 - 1/1000. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.


靶标

功能	Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate.
通路	Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1/5.
序列相似性	Belongs to the glyceraldehyde-3-phosphate dehydrogenase family.
翻译后修饰	S-nitrosylation of Cys-152 leads to interaction with SIAH1, followed by translocation to the nucleus. ISGylated.

细胞定位

Cytoplasm > cytosol. Nucleus. Cytoplasm > perinuclear region. Membrane. Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions.

图片




Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915) at 1/20000 dilution (Purified) + COS-1 (African green monkey kidney fibroblast-like) whole cell lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 36 kDa

Observed band size: 36 kDa




Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915) at 1/20000 dilution (Purified) + HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 36 kDa

Observed band size: 36 kDa



All lanes : Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915) at 1/10000 dilution (unpurified)

Lane 1 : HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) cell lysate

Lane 2 : HeLa (human epithelial cell line from cervix adenocarcinoma) cell lysate

Lane 3 : HepG2 (human liver hepatocellular carcinoma cell line) cell lysate

Lane 4 : HUVEC (human umbilical vein endothelial cell line) cell lysate

Lane 5 : MCF7 (human breast adenocarcinoma cell line) cell lysate

Lane 6 : SH-SY5Y (human neuroblastoma cell line from bone marrow) cell lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : HRP labelled Goat anti-Rabbit IgG at 1/2000 dilution

Predicted band size: 36 kDa


Observed band size: 35 kDa

Secondary antibody - anti-rabbit HRP (ab6721)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GAPDH antibody
[EPR6256] - Loading Control (ab128915)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human bladder carcinoma tissue sections labeling GAPDH with purified ab128915 at 1/2000 dilution (0.06 µg/ml). Perform heat mediated antigen retrieval using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR6256] - Loading Control
(ab128915)

Immunocytochemistry/ Immunofluorescence analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling GAPDH with purified ab128915 at 1/250 dilution (0.4 µg/ml). Cells were fixed in 100% Methanol. Cells were counterstained with [ab195889](#) Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 (2.5 µg/ml) dilution. Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)



Immunoprecipitation - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling GAPDH with purified ab128915 at 1/20 dilution (10µg/ml) (red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).

ab128915 (purified) at 1/20 dilution (0.5ug) immunoprecipitating GAPDH in HeLa whole cell lysates.


Lane 1: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates 10ug

Lane 2 (+): ab128915 & HeLa whole cell lysates

Lane 3 (-): Rabbit monoclonal IgG ([ab172730](#)) instead of ab128915 in HeLa whole cell lysates

For western blotting, VeriBlot for IP Detection Reagent (HRP) ([ab131366](#)) was used at 1/1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)

IHC image of ab128915 (unpurified) staining GAPDH in human pancreas* formalin fixed paraffin embedded tissue sections, performed on a Leica Bond. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab128915, 1:250 dilution, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. No primary antibody was used in the secondary only control (shown on the inset).

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.


*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre



Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)


ab128915 (unpurified) staining GAPDH in HeLa (human epithelial cell line from cervix adenocarcinoma) cells. The cells were fixed with 100% methanol (5min) and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated with ab128915 at 2µg/ml and **ab7291** at 1µg/ml overnight at +4°C, followed by a further incubation at room temperature for 1h with an goat **anti-rabbit AlexaFluor®488 (ab150081)** at 2 µg/ml (shown in green) and goat **anti-mouse AlexaFluor®594 (ab150120)** at 2 µg/ml (shown in pseudo color red). Nuclear DNA was labelled in blue with DAPI.

Negative controls: 1– Rabbit primary antibody and anti-mouse secondary antibody; 2 – Mouse primary antibody and anti-rabbit secondary antibody. Controls 1 and 2 indicate that there is no unspecific reaction between primary and secondary antibodies used.



Flow Cytometry (Intracellular) - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)


Overlay histogram showing HeLa (human epithelial cell line from cervix adenocarcinoma) cells stained with ab128915 (unpurified) (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab128915, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was goat anti-rabbit Alexa Fluor® 488 (H&L) ([ab150077](#)) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.



Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)

This image is courtesy of an Abreview submitted by Kirk McManus

ab128915 (unpurified) staining GAPDH in human HeLa (human epithelial cell line from cervix adenocarcinoma) cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with paraformaldehyde and permeabilized with 0.5% Triton X-100 in PBS. Samples were incubated with primary antibody (1/500 in PBS) for 1 hour at 22°C. An Alexa Fluor® 488-conjugated goat anti-rabbit IgG polyclonal (1/200) was used as the secondary antibody. Counter stained with DAPI.



Immunocytochemistry/ Immunofluorescence - Anti-GAPDH antibody [EPR6256] - Loading Control (ab128915)

ab128915 (unpurified), at 1/250, staining GAPDH in MCF7 (human breast adenocarcinoma cell line) cells by immunofluorescence.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-GAPDH antibody [EPR6256] - Loading Control
(ab128915)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors