abcam

Product datasheet

Anti-Gelsolin antibody [GS-2C4] ab11081



★★★★★ 6 Abreviews 20 References 4 图像

概述

产**品名称** Anti-Gelsolin抗体[GS-2C4]

小鼠单克隆抗体[GS-2C4] to Gelsolin

宿主 Mouse

经测试应用 适用于: WB, ICC/IF, IP

种属反应性 与反应: Human

免疫原 Full length native protein (purified) corresponding to Human Gelsolin.

Database link: P06396-1

表位 The antibody is directed against an epitope located on the 47 kD peptide derived from a

chymotryptic cleavage of gelsolin. This peptide contains the carboxy terminal actin binding site for

gelsolin.

阳性对照 IP: U-2 OS cell lysate. WB: HeLa cell lysate.

常规说明 This product was changed from ascites to tissue culture supernatant on 17 May 2019. Please

note that the dilutions may need to be adjusted accordingly. If you have any questions, please do

not hesitate to contact our scientific support team.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

cycles.

存储溶液 Preservative: 0.097% Sodium azide

纯**度** Tissue culture supernatant

1

 克隆
 单克隆

 克隆编号
 GS-2C4

 同种型
 IgG1

应用


The Abpromise guarantee Abpromise™承诺保证使用ab11081于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	****(4)	Use at an assay dependent concentration. Predicted molecular weight: 86 kDa.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.

靶标	
功能	Calcium-regulated, actin-modulating protein that binds to the plus (or barbed) ends of actin monomers or filaments, preventing monomer exchange (end-blocking or capping). It can promote the assembly of monomers into filaments (nucleation) as well as sever filaments already formed. Plays a role in ciliogenesis.
组织 特异性	Phagocytic cells, platelets, fibroblasts, nonmuscle cells, smooth and skeletal muscle cells.
疾病相关	Defects in GSN are the cause of amyloidosis type 5 (AMYL5) [MIM:105120]; also known as familial amyloidosis Finnish type. AMYL5 is a hereditary generalized amyloidosis due to gelsolin amyloid deposition. It is typically characterized by cranial neuropathy and lattice corneal dystrophy. Most patients have modest involvement of internal organs, but severe systemic disease can develop in some individuals causing peripheral polyneuropathy, amyloid cardiomyopathy, and nephrotic syndrome leading to renal failure.
序列相似性	Belongs to the villin/gelsolin family. Contains 6 gelsolin-like repeats.
翻译后修饰	Phosphorylation on Tyr-86, Tyr-409, Tyr-465, Tyr-603 and Tyr-651 in vitro is induced in presence of phospholipids.
细胞定位	Cytoplasm > cytoskeleton and Secreted.

图片



(ab11081)

All lanes : Anti-Gelsolin antibody [GS-2C4] (ab11081) at 1/200 dilution

Lane 1: Wild-type HeLa cell lysate


Lane 2: GSN knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

Predicted band size: 86 kDa


ab11081 was shown to react with GSN in wild-type HeLa cells in Western blot with loss of signal observed in GSN knockout cell line ab265201 (GSN knockout cell lysate ab257204). Wild-type HeLa and GSN knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab11081 overnight at 4 °C at a 1/200 dilution. Blots were incubated with goat anti-mouse HRP secondary antibodies at 1/5000 before imaging. This data was kindly provided by the YCharOS Inc., an open science company with the mission of characterizing every commercially available antibody reagent. Abcam are working with YCharOS to support their mission of antibody characterisation using knockout cell lines.



Immunocytochemistry/ Immunofluorescence - Anti-Gelsolin antibody [GS-2C4] (ab11081)


ICC/IF image of ab11081 stained HeLa cells. The cells were 4% PFA fixed (10 min) and then incubated in 1% BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab11081, 1 µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-mouse IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue). This antibody also gave a positive IF result in Hek293, HepG2 and MCF7 cells.

This image was generated using the ascites version of the product.



Immunoprecipitation - Anti-Gelsolin antibody [GS-2C4] (ab11081)

Immunoprecipitation of GSN in U-2 OS cells. Lysates were prepared and immunoprecipitation was performed using 1.0 µg of ab11081 pre-coupled to beads. Samples were washed and processed for western blot with ab11081. SM=10% starting material; UB=10% unbound fraction; IP=immunoprecipitate. These data were provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.



Western blot - Anti-Gelsolin antibody [GS-2C4] (ab11081)

Anti-Gelsolin antibody [GS-2C4] (ab11081) at 1/2000 dilution + Human Plasma

Secondary

AP conjugated anti-mouse Fab lgG

Performed under reducing conditions.

Predicted band size: 86 kDa

This image was generated using the ascites version of the product.

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors