

Anti-GFP antibody ab13970

★★★★★ [93 Abreviews](#) [3182 References](#) [8 图像](#)

概述

产品名称	Anti-GFP抗体
描述	鸡多克隆抗体to GFP
宿主	Chicken
特异性	Our GFP antibody does cross-react with the many fluorescent proteins that are derived from the jellyfish <i>Aequorea victoria</i> . These are all proteins that differ from the original GFP by just a few point mutations (EGFP, YFP, mVenus, CFP, BFP etc.).
经测试应用	适用于: WB, ICC/IF
种属反应性	与反应: Species independent
免疫原	Recombinant full length protein corresponding to GFP. Database link: P42212
阳性对照	ICC: GFP-transfected NIH/3T3 (Mouse embryo fibroblast cell line). WB: Transgenic mouse spinal cords.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7 Preservative: 0.01% Thimerosal (merthiolate) Constituents: PBS, 50% Glycerol, 0.16% Sodium phosphate
纯度	IgY fraction
纯化说明	Sterile filtered.
克隆	多克隆

同种型

IgY

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab13970于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (13)	1/5000.
ICC/IF	★★★★★ (22)	1/2000. Used at a dilution of 1/2000 for 1 hr (see Abreview for further information).

靶标

相关性

Function: Energy-transfer acceptor. Its role is to transduce the blue chemiluminescence of the protein aequorin into green fluorescent light by energy transfer. Fluoresces in vivo upon receiving energy from the Ca^{2+} -activated photoprotein aequorin.

Subunit structure: Monomer.


Tissue specificity: Photocytes.

Post-translational modification: Contains a chromophore consisting of modified amino acid residues. The chromophore is formed by autocatalytic backbone condensation between Ser-65 and Gly-67, and oxidation of Tyr-66 to didehydrotyrosine. Maturation of the chromophore requires nothing other than molecular oxygen.

Biotechnological use: Green fluorescent protein has been engineered to produce a vast number of variously colored mutants, fusion proteins, and biosensors. Fluorescent proteins and its mutated allelic forms, blue, cyan and yellow have become a useful and ubiquitous tool for making chimeric proteins, where they function as a fluorescent protein tag. Typically they tolerate N- and C-terminal fusion to a broad variety of proteins. They have been expressed in most known cell types and are used as a noninvasive fluorescent marker in living cells and organisms. They enable a wide range of applications where they have functioned as a cell lineage tracer, reporter of gene expression, or as a measure of protein-protein interactions. Can also be used as a molecular thermometer, allowing accurate temperature measurements in fluids. The measurement process relies on the detection of the blinking of GFP using fluorescence correlation spectroscopy.

Sequence similarities: Belongs to the GFP family.

Biophysicochemical properties: Absorption: Abs(max)=395 nm
Exhibits a smaller absorbance peak at 470 nm. The fluorescence emission spectrum peaks at 509 nm with a shoulder at 540 nm.



All lanes : Anti-GFP antibody (ab13970) at 1/2000 dilution (Diluent 1x TBS /4 hours at 4°C)

Lane 1 : 3 µg of GFP plasmid overexpressed in mouse cardiomyocytes whole cell lysate with BSA / for 1 hour at room temperature

Lane 2 : 2 µg of GFP plasmid overexpressed in mouse cardiomyocytes whole cell lysate with BSA / for 1 hour at room temperature

Lane 3 : 1 µg of GFP plasmid overexpressed in mouse cardiomyocytes whole cell lysate with BSA / for 1 hour at room temperature

Lysates/proteins at 25 µg per lane.

Blocking peptides at 5 % per lane.

Secondary

All lanes : Goat Anti-Chicken IgY H&L (Alexa Fluor® 594) preadsorbed ([ab150176](#)) at 1/5000 dilution


Performed under reducing conditions.

Additional bands at: 25 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 30 seconds

Gel Running Conditions: Reduced Denaturing (15% PAGE)

Detection method: Fluorescent Secondary Antibodies




Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody (ab13970)

This image is courtesy of an Abreview submitted by Christophe Lachaud

ab13970 staining GFP in U-2 OS (Human bone osteosarcoma epithelial cell line) cells by ICC/IF.

Cells were paraformaldehyde fixed, permeabilized with 0.5% triton and blocked with 2% antibody dilution buffer for 2 hours. Cells were incubated with the primary antibody (1/1000) for 1 hour at 25°C. An undiluted Alexa Fluor® 488 conjugated Goat anti-chicken polyclonal was used as the secondary antibody.



Western blot - Anti-GFP antibody (ab13970)

Image courtesy of an anonymous Abreview.

All lanes : Anti-GFP antibody (ab13970) at 1/1000 dilution


Lane 1 : Whole cell lysate prepared from HeLa (Human epithelial cell line from cervix adenocarcinoma) cells.

Lane 2 : Whole cell lysate prepared from HeLa cells.

Lysates/proteins at 25 µg per lane.

Secondary


All lanes : IRDye 800CW conjugated goat anti-chicken polyclonal at 1/15000 dilution



Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody (ab13970)

Transgenic mice expressing GFP selectively in lamina II of the spinal cord.

In the right panels, note the correspondance between the green (rabbit anti-GFP) and red signals (chicken anti-GFP from Abcam) indicating that these two antibody preparations recognized the same gene product. The secondary antibody used with ab13970 was an FITC-labeled goat anti-chicken



Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody (ab13970)


ab13970 staining GFP in GFP-transfected NIH/3T3 (Mouse embryo fibroblast cell line) cells.

The cells were fixed with 4% formaldehyde (10 minutes) and then blocked in 1% BSA / 0.3M glycine in 0.1%PBS-Tween for 1 hour. The cells were then incubated with ab13970 at 1/2000 dilution overnight at +4°C followed by incubation with Goat Anti-Chicken IgY H&L (Alexa Fluor® 488) preadsorbed (**ab150173**), for 1 hour, at 1 µg/ml.

Under identical experimental conditions, when compared to the basal level of GFP expression in transfected NIH/3T3 cells, the cells upon which ab13970 was applied gave a stronger signal in the 488 channel, indicating that ab13970 is binding to GFP and therefore eliciting signal amplification.

ab13970 was also applied to non-GFP-transfected NIH/3T3 cells, which produced no positive staining, indicating specificity for GFP.

Nuclear DNA was labeled with 1.43 µM DAPI (blue).




Western blot - Anti-GFP antibody (ab13970)

Lane 1 : Rabbit anti-GFP

Lane 2 : Anti-GFP antibody (ab13970)

All lanes : Transgenic mouse spinal cords

Western blot of transgenic mouse spinal cords showing that the rabbit anti-GFP (lane 1) and the chicken anti-GFP (Abcam; lane 2) recognize a band at the same molecular weight.




Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody (ab13970)

This image is courtesy of an Abreview submitted by Dr Radbod Darabi

ab13970 staining GFP + tumor in mouse muscle cells by ICC/IF.

Cells were formaldehyde fixed and blocked with 3% BSA for 1 hour at 24°C prior to incubation with the primary antibody (1/500) for 1 hour at 24°C. An Alexa Fluor® 488 conjugated goat anti-chicken was used as the secondary.



Immunocytochemistry/ Immunofluorescence - Anti-GFP antibody (ab13970)

Image courtesy of an AbReview submitted by Dr Francois Daubeuf.

Immunocytochemical/ Immunofluorescence analysis of cytopspined HEK-293 cells (Human epithelial cell line from embryonic kidney) transfected with GFP, labeling GFP with ab13970 at 1/200 incubated for 16 hours at 4°C with 1% BSA in PBS. Secondary used was a donkey anti-chicken polyclonal DyLight® 594 at 1/500.

GFP is shown in red (DyLight® 594).

Nuclei are counterstained in blue (DAPI).

The left panel shows HEK-293 cells transfected with GFP and the right panel shows non-transfected HEK-293 cells.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors