

Anti-GGT1/GGT antibody [EPR5288] ab109427

重组 RabMAb

3 References **5 图像**

概述

产品名称	Anti-GGT1/GGT抗体[EPR5288]
描述	兔单克隆抗体[EPR5288] to GGT1/GGT
宿主	Rabbit
经测试应用	适用于: WB, IHC-P
种属反应性	与反应: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human fetal liver and human fetal kidney tissue lysates and HepG2 and A549 cell lysates. IHC-P: Human kidney and stomach tissues.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.05% BSA</p>
纯度	Protein A purified
克隆	单克隆

克隆编号	EPR5288
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab109427于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000 - 1/10000. Predicted molecular weight: 61 kDa.
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u> .

靶标

功能	Initiates extracellular glutathione (GSH) breakdown, provides cells with a local cysteine supply and contributes to maintain intracellular GSH level. It is part of the cell antioxidant defense mechanism. Catalyzes the transfer of the glutamyl moiety of glutathione to amino acids and dipeptide acceptors. Alternatively, glutathione can be hydrolyzed to give Cys-Gly and gamma glutamate. Isoform 3 seems to be inactive.
组织特异性	Detected in fetal and adult kidney and liver, adult pancreas, stomach, intestine, placenta and lung. Isoform 3 is lung-specific. There are several other tissue-specific forms that arise from alternative promoter usage but that produce the same protein.
通路	Sulfur metabolism; glutathione metabolism.
疾病相关	Defects in GGT1 are a cause of glutathionuria (GLUTH) [MIM:231950]; also known as gamma-glutamyltranspeptidase deficiency. It is an autosomal recessive disease.
序列相似性	Belongs to the gamma-glutamyltransferase family.
翻译后修饰	N-glycosylated on both chains. Contains hexoses, hexosamines and sialic acid residues. Glycosylation profiles tested in kidney and liver tissues reveal the presence of tissue-specific and site-specific glycan composition, despite the overlap in composition among the N-glycans. A total of 36 glycan compositions, with 40 unique structures are observed. Up to 15 different glycans are observed at a single site, with site-specific variation in glycan composition. The difference in glycosylation profiles in the 2 tissues do not affect the enzyme activity.
细胞定位	Membrane.

图片



All lanes : Anti-GGT1/GGT antibody [EPR5288] (ab109427) at 1/1000 dilution (purified)

Lane 1 : Human fetal liver tissue lysate

Lane 2 : HepG2 whole cell lysate

Lane 3 : Human fetal kidney tissue lysate

Lane 4 : A549 whole cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 61 kDa

Observed band size: 50-60 kDa


Blocking and dilution buffer: 5% NFDM/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human stomach tissue labelling GGT1/GGT with purified ab109427 at a dilution of 1/100. Heat mediated antigen retrieval was performed using EDTA buffer pH 9.

ab97051, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Negative control using PBS instead of primary antibody. Counterstained with hematoxylin.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GGT1/GGT antibody [EPR5288] (ab109427)



All lanes : Anti-GGT1/GGT antibody [EPR5288] (ab109427) at 1/1000 dilution (unpurified)

Lane 1 : Human fetal liver lysate

Lane 2 : HepG2 cell lysate

Lane 3 : Human fetal kidney lysate

Lane 4 : A549 cell lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 61 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue labelling GGT1/GGT with unpurified ab109427 at a dilution of 1/100.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?

Research with confidence
Consistent and reproducible results

Long-term and scalable supply
Recombinant technology

Success from the first experiment
Confirmed specificity

Ethical standards compliant
Animal-free production

Anti-GGT1/GGT antibody [EPR5288] (ab109427)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors