


Anti-gp78 antibody ab76841

★★★★★ 1 Abreviews 2 References

概述

产品名称	Anti-gp78抗体
描述	兔多克隆抗体to gp78
宿主	Rabbit
特异性	ab76841 reacts specifically with the 73 kda (isoform 2) protein.
经测试应用	适用于: IHC-Fr, IHC-P, WB
种属反应性	与反应: Human 预测可用于: Mouse 
免疫原	Synthetic peptide corresponding to Human gp78 (C terminal).
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Constituent: Whole serum
纯度	Whole antiserum
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee      **Abpromise™**承诺保证使用ab76841于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr		Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration.
WB	★★★★☆ (1)	Use at an assay dependent concentration.

## 靶标

功能	E3 ubiquitin-protein ligase which can target both itself and other proteins including CD3D and APOB for proteasomal degradation. Mediates polyubiquitination of CYP3A4. Specific receptor for the autocrine motility factor. Mediates tumor invasion and metastasis. Component of a complex required to couple deglycosylation and proteasome-mediated degradation of misfolded proteins in the endoplasmic reticulum that are retrotranslocated in the cytosol. Promotes ubiquitination of misfolded proteins such as mutant CFTR; proposed to mediate sequential ubiquitination by recognizing already ubiquitin-conjugated substrates and to cooperate with E3 ubiquitin-protein ligase RNF5.
通路	Protein modification; protein ubiquitination.
序列相似性	Contains 1 CUE domain. Contains 1 RING-type zinc finger.
细胞定位	Endoplasmic reticulum membrane.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors