

Anti-Heme Oxygenase 1 antibody ab13243

★★★★★ [7 Abreviews](#) [235 References](#) [7 图像](#)

概述

| | |
|-------|---|
| 产品名称 | Anti-Heme Oxygenase 1抗体 |
| 描述 | 兔多克隆抗体to Heme Oxygenase 1 |
| 宿主 | Rabbit |
| 经测试应用 | 适用于: WB, ICC/IF, Sandwich ELISA, IHC-P |
| 种属反应性 | 与反应: Mouse, Rat, Human |
| 免疫原 | Recombinant rat HO-1 (Hsp32) lacking the membrane spanning region |
| 常规说明 | <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p> |

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| 存储溶液 | Preservative: 0.09% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine) |
| 纯度 | Protein A purified |
| 克隆 | 多克隆 |
| 同种型 | IgG |

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab13243于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

| 应用 | Ab评论 | 说明 |
|----------------|-----------|---|
| WB | ★★★★★ (3) | 1/2000. Detects a band of approximately 32 kDa (predicted molecular weight: 34.6 kDa). |
| ICC/IF | | Use a concentration of 1 - 5 µg/ml. |
| Sandwich ELISA | | Use a concentration of 0.5 µg/ml. Can be paired for Sandwich ELISA with Mouse monoclonal [HO-1-1] to Heme Oxygenase 1 (ab13248) . For sandwich ELISA, use this antibody as Detection at 0.5 µg/ml with Mouse monoclonal [HO-1-1] to Heme Oxygenase 1 (ab13248) as Capture. |
| IHC-P | ★★★★★ (3) | Use a concentration of 1 - 5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. |


靶标

| | |
|-------|--|
| 功能 | Heme oxygenase cleaves the heme ring at the alpha methene bridge to form biliverdin. Biliverdin is subsequently converted to bilirubin by biliverdin reductase. Under physiological conditions, the activity of heme oxygenase is highest in the spleen, where senescent erythrocytes are sequestered and destroyed. |
| 序列相似性 | Belongs to the heme oxygenase family. |
| 细胞定位 | Microsome. Endoplasmic reticulum. |

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody (ab13243)



All lanes : Anti-Heme Oxygenase 1 antibody (ab13243) at 1/1000 dilution

Lane 1 : Hek293

Lane 2 : HL60

Lane 3 : HeLa

Lane 4 : A549

Lane 5 : Hu spleen

Lane 6 : Ms spleen

Lane 7 : Rt spleen

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : IRDye® 800CW Goat anti Rabbit

Predicted band size: 34.6 kDa

Observed band size: 32 kDa

Hek293 & HL60 presumed negative or very low expression.


Loading control GAPDH at 38kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody (ab13243)

IHC image of Heme Oxygenase 1 staining in formalin fixed, paraffin embedded rat spleen normal tissue section*, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab13243, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Western blot - Anti-Heme Oxygenase 1 antibody (ab13243)


Anti-Heme Oxygenase 1 antibody (ab13243) (Rat microsomes)

Predicted band size: 34.6 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Heme Oxygenase 1 antibody (ab13243)

Alzheimer diseased section stained with ab13243.



Immunocytochemistry/ Immunofluorescence - Anti-Heme Oxygenase 1 antibody (ab13243)

ICC/IF image of ab13243 stained HepG2 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab13243, 1µg/ml) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



Standard Curve for Heme Oxygenase 1 (Analyte: **Heme Oxygenase 1 protein (Tagged) (ab85243)**); dilution range 1pg/ml to 1µg/ml using Capture Antibody **Mouse monoclonal [HO-1-1] to Heme Oxygenase 1 (ab13248)** at 5µg/ml and Detector Antibody **Rabbit polyclonal to Heme Oxygenase 1 (ab13243)** at 0.5µg/ml.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors