

Anti-HEXIM1 antibody [2416C2a] ab58624

1 图像

概述

产品名称	Anti-HEXIM1抗体[2416C2a]
描述	小鼠单克隆抗体[2416C2a] to HEXIM1
宿主	Mouse
经测试应用	适用于: WB
种属反应性	与反应: Recombinant fragment
免疫原	Recombinant fragment of human HEXIM1
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	pH: 7.40 Preservative: 0.05% Sodium azide Constituents: PBS, 0.0225% Potassium chloride, 0.03% Potassium phosphate, 0.1312% Sodium phosphate, 0.812% Sodium chloride, 1% BSA
纯度	Protein G purified
纯化说明	Purified using protein G column chromatography from culture supernatant of hybridoma cultured in a medium containing bovine IgG-depleted (approximately 95%) fetal bovine serum and filtered through a 0.22µm membrane.
克隆	单克隆
克隆编号	2416C2a
同种型	IgG1

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab58624于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		Use at an assay dependent concentration. Predicted molecular weight: 41 kDa.

靶标

功能	Transcriptional regulator which functions as a general RNA polymerase II transcription inhibitor. In cooperation with 7SK snRNA sequesters P-TEFb in a large inactive 7SK snRNP complex preventing RNA polymerase II phosphorylation and subsequent transcriptional elongation. May also regulate NF-kappa-B, ESR1, NR3C1 and CIITA-dependent transcriptional activity.
组织特异性	Ubiquitously expressed with higher expression in placenta. HEXIM1 and HEXIM2 are differentially expressed. Expressed in endocrine tissues.
序列相似性	Belongs to the HEXIM family.
结构域	The coiled-coil domain mediates oligomerization.
细胞定位	Nucleus. Cytoplasm. Binds alpha-importin and is mostly nuclear.

图片



Anti-HEXIM1 antibody [2416C2a] (ab58624) + immunogen (recombinant pfragment)

Predicted band size: 41 kDa

The MW is low because the recombinant fragment used as immunogen, rather than the full length protein, was used as positive control

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors