


# Anti-hHR23b antibody ab86781

[3 References](#) [5 图像](#)

### 概述

产品名称	Anti-hHR23b抗体
描述	兔多克隆抗体to hHR23b
宿主	Rabbit
经测试应用	适用于: WB, IP, IHC-P
种属反应性	与反应: Mouse, Rat, Human 预测可用于: Rabbit, Horse, Chicken, Guinea pig, Cow, Dog, Pig, Chimpanzee, Rhesus monkey, Gorilla, Tilapia, Orangutan, Platypus 
免疫原	Synthetic peptide corresponding to Human hHR23b aa 350-450. Database link: <a href="#">P54727</a>
阳性对照	HeLa whole cell lysate ( <a href="#">ab150035</a> ) NIH3T3 whole cell lysate
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 0.1% BSA, Tris buffered saline
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

## The Abpromise guarantee

**Abpromise™** 承诺保证使用 ab86781 于以下的经测试应用

“应用说明”部分下显示的仅为推荐的起始稀释度；实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/2000 - 1/10000. Predicted molecular weight: 43 kDa.
IP		Use at 10 µg/mg of lysate.
IHC-P		1/500 - 1/2000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## 靶标

### 功能

Multiubiquitin chain receptor involved in modulation of proteasomal degradation. Binds to polyubiquitin chains. Proposed to be capable to bind simultaneously to the 26S proteasome and to polyubiquitinated substrates and to deliver ubiquitinated proteins to the proteasome. May play a role in endoplasmatic reticulum-associated degradation (ERAD) of misfolded glycoproteins by association with PNGase and delivering deglycosylated proteins to the proteasome. Involved in global genome nucleotide excision repair (GG-NER) by acting as component of the XPC complex. Cooperatively with CETN2 appears to stabilize XPC. May protect XPC from proteasomal degradation.

The XPC complex is proposed to represent the first factor bound at the sites of DNA damage and together with other core recognition factors, XPA, RPA and the TFIIH complex, is part of the pre-precision (or initial recognition) complex. The XPC complex recognizes a wide spectrum of damaged DNA characterized by distortions of the DNA helix such as single-stranded loops, mismatched bubbles or single stranded overhangs. The orientation of XPC complex binding appears to be crucial for inducing a productive NER. XPC complex is proposed to recognize and to interact with unpaired bases on the undamaged DNA strand which is followed by recruitment of the TFIIH complex and subsequent scanning for lesions in the opposite strand in a 5'-to-3' direction by the NER machinery. Cyclobutane pyrimidine dimers (CPDs) which are formed upon UV-induced DNA damage escape detection by the XPC complex due to a low degree of structural perurbation. Instead they are detected by the UV-DDB complex which in turn recruits and cooperates with the XPC complex in the respective DNA repair. In vitro, the XPC:RAD23B dimer is sufficient to initiate NER; it preferentially binds to cisplatin and UV-damaged double-stranded DNA and also binds to a variety of chemically and structurally diverse DNA adducts. XPC:RAD23B contacts DNA both 5' and 3' of a cisplatin lesion with a preference for the 5' side. XPC:RAD23B induces a bend in DNA upon binding. XPC:RAD23B stimulates the activity of DNA glycosylases TDG and SMUG1.

### 序列相似性


Belongs to the RAD23 family.  
Contains 1 ST11 domain.  
Contains 2 UBA domains.  
Contains 1 ubiquitin-like domain.

### 结构域

The ubiquitin-like domain mediates interaction with ATXN3.

### 细胞定位

Nucleus. Cytoplasm. The intracellular distribution is cell cycle dependent. Localized to the nucleus and the cytoplasm during G1 phase. Nuclear levels decrease during S-phase; upon entering mitosis, relocalizes in the cytoplasm without association with chromatin.



Western blot - Anti-hHR23b antibody (ab86781)

**All lanes :** Anti-hHR23b antibody (ab86781) at 0.04  $\mu\text{g/ml}$

**Lane 1 :** HeLa whole cell lysate at 50  $\mu\text{g}$

**Lane 2 :** HeLa whole cell lysate at 15  $\mu\text{g}$


**Lane 3 :** HeLa whole cell lysate at 5  $\mu\text{g}$

**Lane 4 :** NIH3T3 whole cell lysate at 50  $\mu\text{g}$

Developed using the ECL technique.


**Predicted band size:** 43 kDa

**Exposure time:** 30 seconds




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hHR23b antibody (ab86781)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human breast carcinoma tissue labelling hHR23b with ab86781 at 1/1000 (0.2  $\mu\text{g/ml}$ ). Detection: DAB.



Immunoprecipitation - Anti-hHR23b antibody (ab86781)

1 mg HeLa whole cell lysate was immunoprecipitated with 10  $\mu\text{g}$  ab86781 (lane 1) or control IgG (lane 2). 20% of the immunoprecipitate was subjected to Western blotting and labelled with ab86781 at 0.4  $\mu\text{g/ml}$ . Bands were detected by chemiluminescence with an exposure time of 3 seconds.




Western blot - Anti-hHR23b antibody (ab86781)

Anti-hHR23b antibody (ab86781) at 0.2 µg/ml + C6 (Rat glial tumor cell line) whole cell lysate at 50 µg

**Predicted band size:** 43 kDa

**Exposure time:** 3 minutes



Western blot - Anti-hHR23b antibody (ab86781)

**All lanes :** Anti-hHR23b antibody (ab86781) at 0.2 µg/ml

**Lane 1 :** NIH3T3 whole cell lysate

**Lane 2 :** TCMK-1 whole cell lysate

**Lane 3 :** 4T1 whole cell lysate

**Lane 4 :** CT26.WT whole cell lysate

Lysates/proteins at 50 µg per lane.

**Predicted band size:** 43 kDa

**Exposure time:** 30 seconds

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors