

Anti-HHV8 antibody [LN53] ab4103

14 References **1 图像**

概述

产品名称	Anti-HHV8抗体[LN53]
描述	大鼠单克隆抗体[LN53] to HHV8
宿主	Rat
特异性	HHV 8 is the likely etiological agent of Kaposi's sarcoma (KS). HHV 8 encodes a latent nuclear antigen (LNA), which is the product of the viral gene orf 73. LNA is capable of forming a complex with retinoblastoma susceptibility gene product, which may be related to its oncogenic activity.
经测试应用	适用于: IHC-P, IHC-Fr
种属反应性	与反应: Human herpesvirus 8
免疫原	Recombinant full length protein corresponding to HHV8. Corresponding to the latent nuclear antigen 1 molecule of HHV 8.
表位	Located within the ORF-73 region.
阳性对照	Kaposi's sarcoma.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.3</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituents: Tissue culture supernatant, 1% BSA</p>
纯度	IgG fraction
克隆	单克隆

克隆编号LN53

同种型IgG2c

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab4103于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。


应用	Ab评论	说明
IHC-P		1/25 - 1/50. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. ABC method.
IHC-Fr		1/25 - 1/50. ABC method.

靶标

相关性

HHV8 has been found to be associated with three different diseases observed in AIDS patients; kaposi's sarcoma, primary effusion lymphoma (which is a rare type of non-Hodgkin lymphoma affecting the body cavities) and multicentric Castleman's disease. To date there is much evidence to support a direct role for HHV8 in kaposi's sarcoma.

图片



Cultured cells infected with HHV8, stained with HHV8 using ABC and DAB chromogen.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors