

Anti-HIF-2-alpha antibody [ep190b] ab8365

★★★★★ [5 Abreviews](#) [71 References](#) [2 图像](#)

概述

产品名称	Anti-HIF-2-alpha抗体[ep190b]
描述	小鼠单克隆抗体[ep190b] to HIF-2-alpha
宿主	Mouse
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: Rat 
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Hypoxic A549 and HeLa cell lysate.
常规说明	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.02% Sodium azide Constituent: PBS
纯度	Protein G purified
克隆	单克隆
克隆编号	ep190b
骨髓瘤	NS1

同种型	IgG1
轻链类型	kappa

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab8365于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (3)	Use a concentration of 1 µg/ml. Predicted molecular weight: 100 kDa. Abcam recommends using a 3% milk block with this product.

靶标

功能	Transcription factor involved in the induction of oxygen regulated genes. Binds to core DNA sequence 5'-[AG]CGTG-3' within the hypoxia response element (HRE) of target gene promoters. Regulates the vascular endothelial growth factor (VEGF) expression and seems to be implicated in the development of blood vessels and the tubular system of lung. May also play a role in the formation of the endothelium that gives rise to the blood brain barrier. Potent activator of the Tie-2 tyrosine kinase expression. Activation seems to require recruitment of transcriptional coactivators such as CREBPB and probably EP300. Interaction with redox regulatory protein APEX seems to activate CTAD.
组织特异性	Expressed in most tissues, with highest levels in placenta, lung and heart. Selectively expressed in endothelial cells.
疾病相关	Defects in EPAS1 are the cause of erythrocytosis familial type 4 (ECYT4) [MIM:611783]. ECYT4 is an autosomal dominant disorder characterized by increased serum red blood cell mass, elevated hemoglobin concentration and hematocrit, and normal platelet and leukocyte counts.
序列相似性	Contains 1 basic helix-loop-helix (bHLH) domain. Contains 1 PAC (PAS-associated C-terminal) domain. Contains 2 PAS (PER-ARNT-SIM) domains.
翻译后修饰	In normoxia, is probably hydroxylated on Pro-405 and Pro-531 by EGLN1/PHD1, EGLN2/PHD2 and/or EGLN3/PHD3. The hydroxylated prolines promote interaction with VHL, initiating rapid ubiquitination and subsequent proteasomal degradation. Under hypoxia, proline hydroxylation is impaired and ubiquitination is attenuated, resulting in stabilization. In normoxia, is hydroxylated on Asn-847 by HIF1AN thus probably abrogating interaction with CREBBP and EP300 and preventing transcriptional activation. Phosphorylated on multiple sites in the CTAD. The iron and 2-oxoglutarate dependent 3-hydroxylation of asparagine is (S) stereospecific within HIF CTAD domains.
细胞定位	Nucleus.

图片



Western blot - Anti-HIF-2-alpha antibody [ep190b] (ab8365)

All lanes : Anti-HIF-2-alpha antibody [ep190b] (ab8365) at 1/500 dilution

Lane 1 : HeLa whole cell lysate

Lane 2 : HeLa treated with 0.5mM DFO whole cell lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : HRP conjugated Goat Anti-Mouse IgG (H+L) at 1/10000 dilution

Developed using the ECL technique.


Performed under reducing conditions.

Predicted band size: 100 kDa

Observed band size: 100 kDa

Exposure time: 20 minutes

This blot was produced using a 4-12% Bis-tris gel under the MOPS buffer system. The gel was run at 200V for 55 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab8365 overnight at 4°C. Antibody binding was detected using a Goat anti-mouse antibody conjugated to HRP, and visualised using ECL development solution [ab133406](#).



HIF-2-alpha detected in hypoxic Human lysate using ab8365. Lane 1: normoxic A549 lysate control, lane 2: hypoxic A549 lysate.

Western blot - Anti-HIF-2-alpha antibody [ep190b]
(ab8365)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors