abcam

Product datasheet

Anti-hnRNP A2B1 antibody ab31645

★★★★★ <u>5 Abreviews</u> <u>21 References</u> 5 图像

概述

产**品名称** Anti-hnRNP A2B1抗体

描述 兔多克隆抗体to hnRNP A2B1

宿主 Rabbit

特异性 The peptide used as the immunogen for this antibody is found within isoform A2 and B1 of the

protein and should thus recognize a doublet around 35/37 kDa. From Jan 2024, QC testing of replenishment batches of this polyclonal changed. All tested and expected application and reactive species combinations are still covered by our Abcam product promise. However, we no longer test all applications. For more information on a specific batch, please contact our Scientific

Support who will be happy to help.

经测试应用 适用于: IHC-P, WB, ICC, IP

种属反应性 与反应: Human

预测可用于: Mouse, Rat, Chicken, Dog, Xenopus laevis

免疫原 Synthetic peptide corresponding to Human hnRNP A2B1 aa 50-150.

(Peptide available as ab31644)

阳性对照 WB: A549, HEK293T, HeLa, NIH 3T3, and MEF1 cell lysates, and mouse testis tissue lysate.

IHC-P: Human colon tissue. ICC: HeLa cells.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle.

存储溶液 pH: 7.40

Preservative: 0.02% Sodium azide

1

Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

纯**度** Immunogen affinity purified

 克隆
 多克隆

 同种型
 IgG

应用


The Abpromise guarantee Abpromise™承诺保证使用ab31645于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P		Use a concentration of 1 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB	★★★★★ (2)	Use a concentration of 1 - 5 μ g/ml. Detects a band of approximately 35, 38 kDa (predicted molecular weight: 35 , 37 kDa).
ICC		Use a concentration of 5 μg/ml.
IP		Use at an assay dependent concentration.

靶标	
功能	Involved with pre-mRNA processing. Forms complexes (ribonucleosomes) with at least 20 other different hnRNP and heterogeneous nuclear RNA in the nucleous.
序列相似性	Contains 2 RRM (RNA recognition motif) domains.
细胞定位	Nucleus > nucleoplasm. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Component of ribonucleosomes. Predominantly nucleoplasmic, however isoform A2 is also found in the cytoplasm of cells in some tissues. Not found in the nucleolus.

图片



Western blot - Anti-hnRNP A2B1 antibody (ab31645)

All lanes: Anti-hnRNP A2B1 antibody (ab31645) at 1/500 dilution

Lane 1: Wild-type HEK293T cell lysate

Lane 2: HNRNPA2B1 knockout HEK293T cell lysate


Lane 3: A549 cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 35, 37 kDa **Observed band size:** 37 kDa

Lanes 1-3: Merged signal (red and green). Green - ab31645 observed at 37 kDa. Red - loading control, <u>ab7291</u> observed at 50 kDa.


ab31645 Anti-hnRNP A2B1 antibody was shown to specifically react with hnRNP A2B1 in wild-type HEK293T cells. Loss of signal was observed when knockout cell line ab266404 (knockout cell lysate ab257224) was used. Wild-type and hnRNP A2B1 knockout samples were subjected to SDS-PAGE. ab31645 and Anti-alpha Tubulin antibody [DM1A] - Loading Control (ab7291) were incubated overnight at 4°C at 1 in 500 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit lgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse lgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.



Immunocytochemistry - Anti-hnRNP A2B1 antibody (ab31645)

ab31645 staining hnRNP A2B1 in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab31645 at 5 µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin - Loading Control. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), preadsorbed at 1/1000 dilution (shown in green) and ab150120, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 594), pre-adsorbed at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.



Western blot - Anti-hnRNP A2B1 antibody (ab31645)

Anti-hnRNP A2B1 antibody (ab31645) at 1 μ g/ml + HeLa (Human epithelial carcinoma cell line) Whole Cell Lysate at 20 μ g


Secondary

Goat polyclonal to Rabbit lgG (Alexa Fluor® 680) at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 35, 37 kDa **Observed band size:** 35,38 kDa

ab31645 is expected to recognize isoform hnrnp A2 and hnrnp B1. The antibody detects bands at approximately 35 and 38 kDa which are of the correct size to correspond to the predicted molecular weight of these isoforms.



Immunoprecipitation - Anti-hnRNP A2B1 antibody (ab31645)


hnRNP A2B1 was immunoprecipitated using 0.5mg Hela whole cell extract, 5µg of Rabbit polyclonal to hnRNP A2B1 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).

The antibody was incubated under agitation with Protein G beads for 10min, Hela whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab31645.

Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) (ab99697).

Band: 35kDa: hnRNP A2B1.



embedded sections) - Anti-hnRNP A2B1 antibody
(ab31645)

IHC image of ab31645 staining in human colon formalin fixed paraffin embedded tissue section, performed on a Leica Bond TM system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab31645, 1 μ g/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors