

Anti-HOXB4 antibody [EP1919Y] ab76093

重组 RabMAb

2 References **8 图像**

概述

产品名称	Anti-HOXB4抗体[EP1919Y]
描述	兔单克隆抗体[EP1919Y] to HOXB4
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide within Human HOXB4 aa 100-200 (internal sequence). The exact sequence is proprietary.
阳性对照	WB: K-562 lysates. IHC-P: human lung carcinoma tissue. ICC/IF: K-562 cells. Flow Cyt (intra): K-562 cells. IP: K-562 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA</p>
纯度	Protein A purified

克隆	单克隆
克隆编号	EP1919Y
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab76093于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/100. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/1000 - 1/10000. Predicted molecular weight: 28 kDa.
IP		1/60. For unpurified use please use at 1/40
IHC-P		1/100 - 1/250. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. See <u>IHC antigen retrieval protocols</u>
ICC/IF		1/100 - 1/250.

靶标


功能	Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.
序列相似性	Belongs to the Antp homeobox family. Deformed subfamily. Contains 1 homeobox DNA-binding domain.
发展阶段	Expressed in whole embryos and fetuses at 5-9 weeks from conception.
细胞定位	Nucleus.

图片




Immunocytochemistry/ Immunofluorescence - Anti-HOXB4 antibody [EP1919Y] (ab76093)

Immunocytochemistry/ Immunofluorescence analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labeling HOXB4 with purified ab76093 at 1:100 dilution (10 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, **ab150077**) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI (blue) was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Immunoprecipitation - Anti-HOXB4 antibody [EP1919Y] (ab76093)

ab76093 (Purified) at 1:60 dilution (2 µg) immunoprecipitating HOXB4 in K-562 whole cell lysate.
 Lane 1 (input): K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate 10 µg
 Lane 2 (+): ab76093 & K-562 whole cell lysate
 Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of ab76093 in K-562 whole cell lysate
 For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used at 1:1000 dilution.
 Blocking and diluting buffer: 5% NFDm/TBST.




Flow Cytometry (Intracellular) - Anti-HOXB4 antibody
[EP1919Y] (ab76093)

Intracellular Flow Cytometry analysis of K-562 (Human chronic myelogenous leukemia lymphoblast) cells labeling HOXB4 with purified ab76093 at 1/100 dilution (10 µg/ml) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, [ab150077](#)) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HOXB4 antibody
[EP1919Y] (ab76093)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung carcinoma tissue sections labeling HOXB4 with purified ab76093 at 1:100 dilution (11.86 µg/ml). Heat mediated antigen retrieval was performed using [ab93684](#) (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-HOXB4 antibody [EP1919Y] (ab76093)


Anti-HOXB4 antibody [EP1919Y] (ab76093) at 1/1000 dilution
(Purified) + K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysates at 15 µg

Secondary

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 28 kDa

Observed band size: 34 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HOXB4 antibody [EP1919Y] (ab76093)





Immunohistochemical analysis of paraffin-embedded human placental tissue, using unpurified ab76093, at 1/100 dilution.
Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-HOXB4 antibody
[EP1919Y] (ab76093)

Overlay histogram showing K562 cells stained with unpurified ab76093 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab76093, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed.

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-HOXB4 antibody [EP1919Y] (ab76093)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

-
- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors