

Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free ab223535

敲除验证
重组
RabMAb

11 图像

概述

产品名称	Anti-HP1 gamma/CBX3抗体[EPR19802] - BSA and Azide free
描述	兔单克隆抗体[EPR19802] to HP1 gamma/CBX3 - BSA and Azide free
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), ICC/IF, IP, WB, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Human brain lysate; HeLa, HepG2, PC-3, C2C12, 4T1, A549, LNCaP, C6 and NIH/3T3 whole cell lysates; Mouse and rat brain lysates. IHC-P: Human testis and bladder cancer tissues; Mouse liver tissue; Rat kidney tissue. ICC/IF: HeLa and NIH/3T3 cells. Flow Cyt (intra): HeLa cells. IP: HeLa whole cell lysate.
常规说明	<p>ab223535 is the carrier-free version of ab217999.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit</p>

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR19802
同种型	IgG

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用 ab223535 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ICC/IF		Use at an assay dependent concentration.
IP		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 21 kDa (predicted molecular weight: 21 kDa).
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

靶标


功能	Seems to be involved in transcriptional silencing in heterochromatin-like complexes. Recognizes and binds histone H3 tails methylated at 'Lys-9', leading to epigenetic repression. May contribute to the association of the heterochromatin with the inner nuclear membrane through its interaction with lamin B receptor (LBR). Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins.
序列相似性	Contains 2 chromo domains.
翻译后修饰	Phosphorylated by PIM1. Phosphorylated during interphase and possibly hyper-phosphorylated

during mitosis.

细胞定位

Nucleus. Associates with euchromatin and is largely excluded from constitutive heterochromatin. May be associated with microtubules and mitotic poles during mitosis.

图片



This WB data was generated using the same anti-HP1 gamma/CBX3 antibody clone [EPR19802] in a different buffer format (cat# **ab217999**).


Lane 1: Wild type HAP1 whole cell lysate (20 µg)

Lane 2: CBX3 knockout HAP1 whole cell lysate (20 µg)

Lane 3: HepG2 whole cell lysate (20 µg)

Lanes 1 - 3: Merged signal (red and green). Green - **ab217999** observed at 25 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab217999 was shown to recognize CBX3 when CBX3 knockout samples were used, along with additional cross-reactive bands. Wild-type and CBX3 knockout samples were subjected to SDS-PAGE. Ab217999 and **ab9484** (Mouse anti GAPDH loading control) were incubated overnight at 4°C at 2000 dilution and 1/10000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunohistochemical analysis of paraffin-embedded human testis tissue labeling HP1 gamma/CBX3 with **ab217999** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution. Nuclear staining on human testis is observed [PMID: 19786570]. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

All lanes : Anti-HP1 gamma/CBX3 antibody [EPR19802] (**ab217999**) at 1/2000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : CBX3 knockout HeLa cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.


Predicted band size: 21 kDa

Observed band size: 25 kDa

This data was developed using the same antibody clone in a different buffer formulation (**ab217999**).

Lanes 1-2: Merged signal (red and green). Green - **ab217999** observed at 25 kDa. Red - Anti-alpha Tubulin antibody [DM1A] - Loading Control (**ab7291**) observed at 50 kDa.

ab217999 was shown to react with HP1 gamma/CBX3 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line **ab261744** (knockout cell lysate **ab257110**) was used. Wild-type HeLa and CBX3 knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. **ab217999** and Anti-alpha Tubulin antibody [DM1A] - Loading Control (**ab7291**) overnight at 4°C at a 1 in 2000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling HP1 gamma/CBX3 with **ab217999** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution. Nuclear staining on hepatocytes of mouse liver is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunohistochemical analysis of paraffin-embedded human bladder cancer tissue labeling HP1 gamma/CBX3 with **ab217999** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution. Nuclear staining on tumor cells of human bladder cancer is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunohistochemical analysis of paraffin-embedded rat kidney tissue labeling HP1 gamma/CBX3 with **ab217999** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution. Nuclear staining on rat kidney is observed. Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/500 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling HP1 gamma/CBX3 with **ab217999** at 1/1000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing nuclear staining on HeLa cell line.

The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).




Immunocytochemistry/ Immunofluorescence - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryonic fibroblast cell line) cells labeling HP1 gamma/CBX3 with **ab217999** at 1/1000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) secondary antibody at 1/1000 dilution (green). Confocal image showing nuclear staining on NIH/3T3 cell line.

The nuclear counterstain is DAPI (blue). Tubulin is detected with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) at 1/1000 dilution.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).



Flow Cytometry (Intracellular) - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling HP1 gamma/CBX3 with **ab217999** at 1/400 dilution (red) compared with a rabbit monoclonal IgG isotype control (**ab172730**; black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (Alexa Fluor[®] 488) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).



Immunoprecipitation - Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

HP1 gamma/CBX3 was immunoprecipitated from 0.35 mg of HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate with **ab217999** at 1/30 dilution. Western blot was performed from the immunoprecipitate using **ab217999** at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (**ab131366**), was used for detection at 1/10000 dilution.

Lane 1: HeLa whole cell lysate, 10 µg (Input).

Lane 2: **ab217999** IP in HeLa whole cell lysate.





Lane 3: Rabbit monoclonal IgG (**ab172730**) instead of **ab217999** in HeLa whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 1 second.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab217999**).

Why choose a recombinant antibody?

 <p>Research with confidence Consistent and reproducible results</p>	 <p>Long-term and scalable supply Recombinant technology</p>
 <p>Success from the first experiment Confirmed specificity</p>	 <p>Ethical standards compliant Animal-free production</p>

Anti-HP1 gamma/CBX3 antibody [EPR19802] - BSA and Azide free (ab223535)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors