

### Anti-HSF1 antibody [EP1710Y] - CHIP Grade ab52757

敲除验证
重组
RabMAb

[26 References](#)
[15 图像](#)

#### 概述

<b>产品名称</b>	Anti-HSF1抗体[EP1710Y] - CHIP Grade
<b>描述</b>	兔单克隆抗体[EP1710Y] to HSF1 - CHIP Grade
<b>宿主</b>	Rabbit
<b>经测试应用</b>	<b>适用于:</b> Flow Cyt (Intra), ChIP, WB, IP, IHC-P, ICC/IF
<b>种属反应性</b>	<b>与反应:</b> Mouse, Human
<b>免疫原</b>	Synthetic peptide within Human HSF1 aa 450 to the C-terminus (C terminal). The exact sequence is proprietary.
<b>阳性对照</b>	WB: K562 HAP1 and HeLa whole cell lysate ( <a href="#">ab150035</a> ). ICC/IF: MCF-7 cells. Flow Cyt (intra): HeLa cells. IHC-P: Human ovarian carcinoma tissue; Mouse testis and colon tissue.
<b>常规说明</b>	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> <li>- High batch-to-batch consistency and reproducibility</li> <li>- Improved sensitivity and specificity</li> <li>- Long-term security of supply</li> <li>- Animal-free production</li> </ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

#### 性能

<b>形式</b>	Liquid
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle. Stable for 12 months at -20°C.
<b>存储溶液</b>	<p>pH: 7.20</p> <p>Preservative: 0.01% Sodium azide</p> <p>Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.21% BSA</p>
<b>纯度</b>	Protein A purified

克隆	单克隆
克隆编号	EP1710Y
同种型	IgG

## 应用

**The Abpromise guarantee** **Abpromise™**承诺保证使用ab52757于以下的经测试应用

“应用说明”部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/20. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
ChIP		Use at an assay dependent concentration.
WB		1/50000. Detects a band of approximately 85 kDa (predicted molecular weight: 57 kDa). <b>For unpurified use at 1/100000.</b>
IP		1/20. <b>For unpurified use at 1/100.</b>
IHC-P		1/250. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		1/100 - 1/250.

## 靶标


功能	DNA-binding protein that specifically binds heat shock promoter elements (HSE) and activates transcription. In higher eukaryotes, HSF is unable to bind to the HSE unless the cells are heat shocked.
序列相似性	Belongs to the HSF family.
结构域	the 9aaTAD motif is a transactivation domain present in a large number of yeast and animal transcription factors.
翻译后修饰	<p>Phosphorylated on multiple serine residues, a subset of which are involved in stress-related regulation of transcription activation. Constitutive phosphorylation represses transcriptional activity at normal temperatures. Levels increase on specific residues heat-shock and enhance HSF1 transactivation activity. Phosphorylation on Ser-307 derepresses activation on heat-stress and in combination with Ser-303 phosphorylation appears to be involved in recovery after heat-stress. Phosphorylated on Ser-230 by CAMK2, in vitro. Cadmium also enhances phosphorylation at this site. Phosphorylation on Ser-303 is a prerequisite for HSF1 sumoylation. Phosphorylation on Ser-121 inhibits transactivation and promotes HSP90 binding. Phosphorylation on Thr-142 also mediates transcriptional activity induced by heat. Phosphorylation on Ser-326 plays an important role in heat activation of HSF1 transcriptional activity.</p> <p>Sumoylated with SUMO1 and SUMO2 on heat-shock. Heat-inducible sumoylation occurs after 15 min of heat-shock, after which levels decrease and at 4 hours, levels return to control levels. Sumoylation has no effect on HSE binding nor on transcriptional activity. Phosphorylation on Ser-</p>

303 is a prerequisite for sumoylation.

## 细胞定位

Cytoplasm. Nucleus. Cytoplasmic during normal growth. On activation, translocates to nuclear stress granules. Colocalizes with SUMO1 in nuclear stress granules.

## 图片



**All lanes :** Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757) at 1/100000 dilution

**Lane 1 :** Wild-type HAP1 whole cell lysate

**Lane 2 :** Hsf1 knockout HAP1 whole cell lysate

**Lane 3 :** HeLa whole cell lysate

**Lane 4 :** K562 whole cell lysate


Lysates/proteins at 20 µg per lane.

**Predicted band size:** 57 kDa

**Observed band size:** 57 kDa


**Lanes 1 - 4:** Merged signal (red and green). Green - ab52757 observed at 57 kDa. Red - loading control, **ab9484**, observed at 37 kDa.

ab52757 was shown to specifically react with in wild-type HAP1 cells as signal was lost in Hsf1 knockout cells. Wild-type and Hsf1 knockout samples were subjected to SDS-PAGE. Ab52757 and **ab9484** (Mouse anti-GAPDH loading control) were incubated overnight at 4°C at 1/100000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human ovarian carcinoma tissue sections labeling HSF1 with Purified ab52757 at 1:250 dilution (1.06 µg/ml). Heat mediated antigen retrieval was performed using **ab93684** (Tris/EDTA buffer, pH 9.0). Tissue was counterstained with Hematoxylin. ImmunoHistoProbe one step HRP Polymer (ready to use) secondary antibody was used at 1:0 dilution. PBS instead of the primary antibody was used as the negative control.



Immunoprecipitation - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

ab52757 (purified) at 1:20 dilution (2µg) immunoprecipitating HSF1 in HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate.


**Lane 1 (input):** HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10ug

**Lane 2 (+):** ab52757 & HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

**Lane 3 (-):** Rabbit monoclonal IgG (**ab172730**) instead of **ab52727** in HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP) (**ab131366**) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDm/TBST.




ChIP - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Chromatin was prepared from HeLa cells heat shocked (42°C 30 minutes) according to the Abcam Dual X-ChIP protocol. Cells were fixed with EGS for 30 minutes, then formaldehyde for 10 minutes. The ChIP was performed with 25 µg of chromatin, 5 µg of ab52757 (red), and 20 µl of Protein A/G sepharose beads. 5 µg of rabbit normal IgG was added to the beads control (gray). The immunoprecipitated DNA was quantified by real time PCR (sybr green approach).

Primers and probes are located in the first kb of the transcribed region.


\*<http://www.abcam.com/resources?>

keywords=X%20ChIP%20protocol




Immunocytochemistry/ Immunofluorescence - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Immunocytochemistry/ Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma epithelial cell) cells labeling HSF1 with Purified ab52757 at 1:100 dilution. Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200. **ab150077** Goat anti rabbit IgG(Alexa Fluor® 488) was used as the secondary antibody at 1:1000 dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling HSF1 with purified ab52757 at 1/20 dilution (10µg/ml) (red). Cells were fixed with 80% Methanol and permeabilised with 0.1% Tween-20. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cell without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757) at 1/50000 dilution (purified) + HeLa (Human cervix adenocarcinoma epithelial cell ) whole cell lysates at 15 µg


**Secondary**

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

**Predicted band size:** 57 kDa

**Observed band size:** 80 kDa

Blocking and diluting buffer: 5% NFDm/TBST



Western blot - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)


Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757) at 1/100000 dilution (unpurified) + HeLa cell lysate at 10 µg

**Secondary**

HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution

**Predicted band size:** 57 kDa

**Observed band size:** 85 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling HSF1 with ab52757 at 1/1000 followed by a Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) Ready to use. Nuclear staining on mouse testis. The section was incubated with ab52757 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND<sup>®</sup> RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**), Ready to use.

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Immunohistochemical analysis of paraffin-embedded Mouse colon tissue labeling HSF1 with ab52757 at 1/1000 followed by a Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) Ready to use. Nuclear staining on mouse colon. The section was incubated with ab52757 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND<sup>®</sup> RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**), Ready to use.


Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

ab52757 staining HSF1 in wild-type Hap1 cells (top panel) and HSF1 knockout Hap1 cells (bottom panel). The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab52757 at 1/250 dilution and **ab7291** (Mouse monoclonal to alpha Tubulin) at 1/1000 dilution overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 µg/ml (shown in green) and a goat secondary antibody to mouse IgG (Alexa Fluor® 594) (**ab150120**) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Immunocytochemistry/Immunofluorescence analysis of MCF-7 cells labelling HSF1 with unpurified ab52757 at 1/1000. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody.

Control: PBS only.

Nuclear counter stain: DAPI.




Flow Cytometry (Intracellular) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

Overlay histogram showing HeLa cells stained with unpurified ab52757 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified ab52757, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10<sup>6</sup> cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10



min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunohistochemical staining of paraffin-embedded human ovarian carcinoma using unpurified ab52757 at a 1:100 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

### Why choose a recombinant antibody?

 <p><b>Research with confidence</b> Consistent and reproducible results</p>	 <p><b>Long-term and scalable supply</b> Recombinant technology</p>
 <p><b>Success from the first experiment</b> Confirmed specificity</p>	 <p><b>Ethical standards compliant</b> Animal-free production</p>

Anti-HSF1 antibody [EP1710Y] - ChIP Grade (ab52757)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors