



### Anti-Hsp27 antibody ab5579

★★★★★ [1 Abreviews](#) [19 References](#) [8 图像](#)

#### 概述

产品名称	Anti-Hsp27抗体
描述	兔多克隆抗体to Hsp27
宿主	Rabbit
经测试应用	适用于: WB, ICC/IF, IHC-P
种属反应性	与反应: Mouse, Rat, Human, African green monkey
免疫原	Synthetic peptide corresponding to Human Hsp27 aa 10-21. Sequence: LLRGPSWDPFRC  (Peptide available as <a href="#">ab39789</a> )
阳性对照	<div>  <a href="#">Run BLAST with</a>  <a href="#">Run BLAST with</a> </div> WB: HEK-293T, HeLa, K562, A431, HepG2, COS-7, NIH/3T3, MCF7, MDA-MB-231, PC3, DU 145, LNCaP, HT1080 whole cell lysate. IHC-P: Human skeletal muscle and breast carcinoma tissue. ICC/IF: HeLa, MCF-7 and C6 cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Sodium azide Constituents: 99% PBS, 0.1% BSA
纯度	Immunogen affinity purified
纯化说明	Antigen affinity chromatography.

克隆

多克隆

同种型

IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab5579于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (1)	1/1000 - 1/2000. Detects a band of approximately 27 kDa.
ICC/IF		1/50.
IHC-P		Use a concentration of 4 µg/ml. Perform heat mediated antigen retrieval via the microwave method before commencing with IHC staining protocol.

靶标

功能

Involved in stress resistance and actin organization.

组织特异性

Detected in all tissues tested: skeletal muscle, heart, aorta, large intestine, small intestine, stomach, esophagus, bladder, adrenal gland, thyroid, pancreas, testis, adipose tissue, kidney, liver, spleen, cerebral cortex, blood serum and cerebrospinal fluid. Highest levels are found in the heart and in tissues composed of striated and smooth muscle.

疾病相关

Defects in HSPB1 are the cause of Charcot-Marie-Tooth disease type 2F (CMT2F) [MIM:606595]. CMT2F is a form of Charcot-Marie-Tooth disease, the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy or CMT1, and primary peripheral axonal neuropathy or CMT2. Neuropathies of the CMT2 group are characterized by signs of axonal regeneration in the absence of obvious myelin alterations, normal or slightly reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy. Nerve conduction velocities are normal or slightly reduced. CMT2F onset is between 15 and 25 years with muscle weakness and atrophy usually beginning in feet and legs (peroneal distribution). Upper limb involvement occurs later. CMT2F inheritance is autosomal dominant.  
Defects in HSPB1 are a cause of distal hereditary motor neuronopathy type 2B (HMN2B) [MIM:608634]. Distal hereditary motor neuronopathies constitute a heterogeneous group of neuromuscular disorders caused by selective impairment of motor neurons in the anterior horn of the spinal cord, without sensory deficit in the posterior horn. The overall clinical picture consists of a classical distal muscular atrophy syndrome in the legs without clinical sensory loss. The disease starts with weakness and wasting of distal muscles of the anterior tibial and peroneal compartments of the legs. Later on, weakness and atrophy may expand to the proximal muscles of the lower limbs and/or to the distal upper limbs.

序列相似性

Belongs to the small heat shock protein (HSP20) family.

翻译后修饰


Phosphorylated in MCF-7 cells on exposure to protein kinase C activators and heat shock.

细胞定位

Cytoplasm. Nucleus. Cytoplasm > cytoskeleton > spindle. Cytoplasmic in interphase cells. Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock

and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles.

## 图片



**All lanes :** Anti-Hsp27 antibody (ab5579) at 1/2000 dilution

**Lane 1 :** MCF7 (human breast adenocarcinoma cell line) whole cell lysate

**Lane 2 :** MDA-MB-231 (human breast adenocarcinoma cell line) whole cell lysate

**Lane 3 :** PC3 (human prostate adenocarcinoma cell line) whole cell lysate

**Lane 4 :** DU 145 (human prostate carcinoma cell line) whole cell lysate

**Lane 5 :** HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate


**Lane 6 :** LNCaP (human prostate cancer cell line) whole cell lysate

**Lane 7 :** HT1080 (human fibrosarcoma cell line) whole cell lysate


Lysates/proteins at 30 µg per lane.

## Secondary

**All lanes :** Goat anti-Rabbit IgG (H+L) Superclonal™ Recombinant Secondary Antibody, HRP at 1/4000 dilution




Immunofluorescence analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Hsp27 (green) with ab5579 at 1/200 dilution, followed by DyLight 488-conjugated secondary antibody. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. 60X magnification. Right - negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp27 antibody (ab5579)


Immunohistochemical analysis of both normal and cancer biopsies of deparaffinized human skeletal muscle tissue labeling Hsp27 with ab5579 at 1/20 dilution or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP, followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.

To expose target proteins, heat induced antigen retrieval was performed using 10mM sodium citrate (pH 6.0) buffer, microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature.



Immunocytochemistry/ Immunofluorescence - Anti-Hsp27 antibody (ab5579)


Immunofluorescence analysis of MCF7 (Human breast adenocarcinoma cell line) cells labeling Hsp27 (green) with ab5579 at 1/200 dilution, followed by DyLight 488-conjugated secondary antibody. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. 60X magnification. Right - negative control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp27 antibody (ab5579)


Immunohistochemical analysis of both normal and cancer biopsies of deparaffinized human breast carcinoma tissue labeling Hsp27 with ab5579 at 1/100 dilution or without primary antibody (negative control) overnight at 4°C in a humidified chamber. Tissues were washed extensively with PBST and endogenous peroxidase activity was quenched with a peroxidase suppressor. Detection was performed using a biotin-conjugated secondary antibody and SA-HRP, followed by colorimetric detection using DAB. Tissues were counterstained with hematoxylin and prepped for mounting.

To expose target proteins, heat induced antigen retrieval was performed using 10mM sodium citrate (pH6.0) buffer, microwaved for 8-15 minutes. Following antigen retrieval tissues were blocked in 3% BSA-PBS for 30 minutes at room temperature.



Immunocytochemistry/ Immunofluorescence - Anti-Hsp27 antibody (ab5579)

Immunofluorescence analysis of C6 (Rat glial tumor cell line) cells labeling Hsp27 (green) with ab5579 at 1/100 dilution, followed by DyLight 488-conjugated secondary antibody. F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue). Cells were fixed with formaldehyde and incubated with the primary antibody overnight at 4°C. 60X magnification. Right - negative control.



Western blot - Anti-Hsp27 antibody (ab5579)

**All lanes :** Anti-Hsp27 antibody (ab5579) at 1/1000 dilution

**Lane 1 :** HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 2 :** HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate

**Lane 3 :** K562 (human chronic myelogenous leukemia lymphoblast cell line ) whole cell lysate

**Lane 4 :** A431 (human epidermoid carcinoma cell line) whole cell lysate

**Lane 5 :** HepG2 (human liver hepatocellular carcinoma cell line) whole cell lysate


**Lane 6 :** COS-7 (african green monkey kidney fibroblast-like cell line) whole cell lysate

**Lane 7 :** NIH/3T3 (mouse embryonic fibroblast cell line) whole cell lysate

Lysates/proteins at 50 µg per lane.

## Secondary

**All lanes :** Goat anti-rabbit IgG-HRP secondary antibody at 1/20000 dilution



Immunocytochemistry/ Immunofluorescence - Anti-Hsp27 antibody (ab5579)

This image is courtesy of Michael Manicini, Ph.D.

HeLa (human epithelial cell line from cervix adenocarcinoma) cells were fixed with 4% formaldehyde in PEM buffer. The coverslip was incubated in blocking buffer of 5% powdered milk in TBS-T plus 0.02% sodium azide for 1 hour at room temperature. Blocking buffer was removed and primary antibody was added at a dilution of 1/250 and incubated overnight at 4 degrees celsius. The coverslips were then washed 4-5 times with blocking buffer for 5 minutes. Secondary antibody, goat anti-rabbit Alexa 594 (**ab150080**), was added at a dilution of 1/1000 and incubated at room temperature for one hour. From this point on coverslips were covered with foil to protect them from light. They were washed 5 times with TBS-T and then one time with PEM, for 5 minutes each wash. The coverslips were fixed 10-30 minutes in 4% formaldehyde in PEM buffer, then washed 3 times with PEM buffer for 5 minutes. 0.1M ammonium chloride in PEM buffer was added for 10 minutes to quench autofluorescence, and then slips were washed 2 times for 5 minutes in PEM followed by 3 washes for 5 minute

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors