

Anti-Hsp27 (phospho S86) antibody ab17938

★★★★★ [2 Abreviews](#) [3 References](#) [2 图像](#)

概述

产品名称	Anti-Hsp27 (phospho S86)抗体
描述	兔多克隆抗体to Hsp27 (phospho S86)
宿主	Rabbit
特异性	Lysates prepared from NIH3T3 cells were immunoblotted in the presence of non-phosphopeptide corresponding to the immunogen, a generic phosphoserine-containing peptide or the phosphopeptide immunogen. The data show that only the peptide corresponding to the mouse phospho S86 HSP27 blocks the antibody signal, thereby demonstrating the specificity of the antibody. The signal was completely removed by lambda phosphatase treatment demonstrating that the antibody interacts specifically with the phosphorylated protein.
经测试应用	适用于: WB, ICC
种属反应性	与反应: Mouse, Human
免疫原	Synthetic peptide corresponding to Mouse Hsp27 (phospho S86).
阳性对照	WB: NIH/3T3 cells. ICC: HeLa cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	pH: 7.30 Preservative: 0.05% Sodium azide Constituents: PBS, 50% Glycerol (glycerin, glycerine), 0.1% BSA
纯度	Immunogen affinity purified
纯化说明	The antibody has been negatively preadsorbed using a non-phosphopeptide corresponding to the

site of phosphorylation to remove antibody that is reactive with non-phosphorylated HSP25 (the mouse homolog of human HSP27). The final product is generated by affinity chromatography using an HSP25-derived peptide that is phosphorylated at serine 86.

克隆 多克隆
同种型 IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab17938于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	★★★★★ (1)	1/1000. Detects a band of approximately 23 kDa (predicted molecular weight: 23 kDa).
ICC		1/250.

靶标

功能 Involved in stress resistance and actin organization.

组织特异性 Detected in all tissues tested: skeletal muscle, heart, aorta, large intestine, small intestine, stomach, esophagus, bladder, adrenal gland, thyroid, pancreas, testis, adipose tissue, kidney, liver, spleen, cerebral cortex, blood serum and cerebrospinal fluid. Highest levels are found in the heart and in tissues composed of striated and smooth muscle.

疾病相关 Defects in HSPB1 are the cause of Charcot-Marie-Tooth disease type 2F (CMT2F) [MIM:606595]. CMT2F is a form of Charcot-Marie-Tooth disease, the most common inherited disorder of the peripheral nervous system. Charcot-Marie-Tooth disease is classified in two main groups on the basis of electrophysiologic properties and histopathology: primary peripheral demyelinating neuropathy or CMT1, and primary peripheral axonal neuropathy or CMT2. Neuropathies of the CMT2 group are characterized by signs of axonal regeneration in the absence of obvious myelin alterations, normal or slightly reduced nerve conduction velocities, and progressive distal muscle weakness and atrophy. Nerve conduction velocities are normal or slightly reduced. CMT2F onset is between 15 and 25 years with muscle weakness and atrophy usually beginning in feet and legs (peroneal distribution). Upper limb involvement occurs later. CMT2F inheritance is autosomal dominant.

Defects in HSPB1 are a cause of distal hereditary motor neuronopathy type 2B (HMN2B) [MIM:608634]. Distal hereditary motor neuronopathies constitute a heterogeneous group of neuromuscular disorders caused by selective impairment of motor neurons in the anterior horn of the spinal cord, without sensory deficit in the posterior horn. The overall clinical picture consists of a classical distal muscular atrophy syndrome in the legs without clinical sensory loss. The disease starts with weakness and wasting of distal muscles of the anterior tibial and peroneal compartments of the legs. Later on, weakness and atrophy may expand to the proximal muscles of the lower limbs and/or to the distal upper limbs.


序列相似性 Belongs to the small heat shock protein (HSP20) family.

翻译后修饰 Phosphorylated in MCF-7 cells on exposure to protein kinase C activators and heat shock.

细胞定位 Cytoplasm. Nucleus. Cytoplasm > cytoskeleton > spindle. Cytoplasmic in interphase cells.

Colocalizes with mitotic spindles in mitotic cells. Translocates to the nucleus during heat shock and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles.

图片



Hsp25 (phospho S86) antibody image 6534.

Western blot using ab17938 on NIH3T3 cells treated with anisomycin.

Lane 1: unstimulated cells

Lane 2: cells stimulated with anisomycin


Lane 3: cells stimulated with anisomycin. Antibody blocked with the non-phosphopeptide corresponding to the immunogen

Lane 4: cells stimulated with anisomycin. Antibody blocked with generic phosphoserine-containing peptide

Lane 5: cells stimulated with anisomycin. Antibody blocked with the phosphopeptide immunogen

Lane 6: cells stimulated with anisomycin and treated with lambda phosphatase

10-30 µg of cell lysate can be loaded when using similar lysates with this antibody. Samples were run using SDS-PAGE on a 10% polyacrylamide gel and transferred to PVDF. Membranes were blocked with a 5% BSA-TBST buffer for one hour at room temperature, then incubated with ab17938 for one hour at room temperature in 3% BSA-TBST buffer, following prior incubation with blocking



HeLa cells stained for Hsp27 (green) using ab17938 at 1/250 dilution in ICC/IF. It was followed by Goat anti-Rabbit IgG (H+L) Superclonal™ Secondary Antibody, Alexa Fluor® 488 conjugate at 1/2000 dilution for 45 minutes at room temperature (Panel a). Nuclei (Panel b: blue) were stained with SlowFade® Gold Antifade Mountant with DAPI. F-actin (Panel c: red) was stained with Rhodamine Phalloidin at 1/300 dilution. Panel d is a merged image showing cytoplasmic localization. Panel e is untreated cell with no signal. Panel f is a no primary antibody control.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors