

Anti-Hsp70 antibody [3A3] ab5439

重组

★★★★★ **3 Abreviews** **57 References** **9 图像**

概述

| | |
|-------|---|
| 产品名称 | Anti-Hsp70抗体[3A3] |
| 描述 | 小鼠单克隆抗体[3A3] to Hsp70 |
| 宿主 | Mouse |
| 特异性 | This antibody detects several members of the heat shock protein 70 kDa (Hsp 70) gene family including Hsp 70, Hsc 70, p75, and following heat shock, Hsp 72 from yeast, Drosophila, fish, mouse, avian, amphibian and human samples. |
| 经测试应用 | 适用于: ICC/IF, IHC-P, WB |
| 种属反应性 | 与反应: Mouse, Rat, Human |
| 免疫原 | Recombinant fragment corresponding to Human Hsp70 aa 504-617 (C terminal). |
| 表位 | Epitope mapping with a panel of Hsp 70 deletion mutants suggests that the epitope recognized is located between amino acids 504-617 of human Hsp 70, a region that has been shown to be involved in stress-induced nucleolar localization. |
| 阳性对照 | WB: HeLa, K562, Jurkat, 293T, A431, A549, F9, RAW264.7 and C6, PC-12 lysates. IHC-P: Human cerebrum and lung carcinoma, Mouse testis, Rat cerebrum tissues. ICC/IF: HeLa and NIH/3T3, PC-12 cells. |
| 常规说明 | <p>This product has switched from a hybridoma to recombinant production method on 08th March 2021.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> |

性能

| | |
|------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. |
| 存储溶液 | Preservative: 0.01% Sodium azide |

| | |
|--------------------|---|
| 纯度 | Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| Primary antibody说明 | Protein A purified The Hsp 70 family is a set of highly conserved proteins that are induced by a variety of biological stresses, including heat stress, in every organism in which the proteins have been examined. The human Hsp 70 family members include: Hsp 70, a protein which is strongly inducible in all organisms but which is also constitutively expressed in primate cells; Hsp 72, a 72 kDa protein that is induced exclusively under stress conditions; Hsc 70, or cognate protein, is a 72 kDa, constitutively expressed, protein which is involved in the uncoating of clathrin coated vesicles; GRP78, or BiP, is a glucose regulated 78 kDa protein localized in the endoplasmic reticulum; and p75, or Hsp 75, a 75 kDa protein that is found within the mitochondria. |
| 克隆 | 单克隆 |
| 克隆编号 | 3A3 |
| 同种型 | IgG1 |

应用

The Abpromise guarantee **Abpromise™** 承诺保证使用ab5439于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度; 实际最佳的稀释度/浓度应由使用者检定。


| 应用 | Ab评论 | 说明 |
|--------|-----------|--|
| ICC/IF | | 1/50. |
| IHC-P | | 1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. |
| WB | ★★★★★ (3) | 1/1000. Predicted molecular weight: 70-78 kDa. |

靶标

相关性 Function: In cooperation with other chaperones, the Hsp70 family stabilize preexistent proteins against aggregation and mediate the folding of newly translated polypeptides in the cytosol as well as within organelles. These chaperones participate in all these processes through their ability to recognize nonnative conformations of other proteins. They bind extended peptide segments with a net hydrophobic character exposed by polypeptides during translation and membrane translocation, or following stress-induced damage. In case of rotavirus A infection, serves as a post-attachment receptor for the virus to facilitate entry into the cell. Tissue specificity: HSPA1B is testis-specific.

细胞定位 Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.

图片



All lanes : Anti-Hsp70 antibody [3A3] (ab5439) at 1/1000 dilution

Lane 1 : HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate

Lane 2 : K562 (human chronic myelogenous leukemia lymphoblast), whole cell lysate

Lane 3 : Jurkat (human T cell leukemia T lymphocyte), whole cell lysate

Lane 4 : 293T (human embryonic kidney epithelial cell), whole cell lysate

Lane 5 : A431 (human epidermoid carcinoma epithelial cell), whole cell lysate

Lane 6 : A549 (human lung carcinoma epithelial cell), whole cell lysate

Lane 7 : F9 (mouse embryonal carcinoma epithelial cell), whole cell lysate

Lane 8 : RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage), whole cell lysate

Lane 9 : C6 (rat glial tumor glial cell), whole cell lysate

Lane 10 : PC-12 (rat adrenal gland pheochromocytoma), whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Peroxidase-Conjugated Goat anti-Mouse IgG (H+L) at 1/10000 dilution

Predicted band size: 70-78 kDa

Observed band size: 70 kDa

Exposure time: 1 second

Blocking/Diluting buffer and concentration: 5% NFDM/TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp70 antibody [3A3] (ab5439)

Immunohistochemical analysis of paraffin-embedded Human cerebrum tissue labeling Hsp70 with ab5439 at 1/2000 (0.519 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human cerebrum. The section was incubated with ab5439 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-Hsp70 antibody [3A3] (ab5439)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized HeLa cells labelling Hsp70 with ab5439 at 1/50 (20.76 µg/ml) dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic and weak nuclear staining in HeLa cells **ab179513** Anti-beta Tubulin rabbit monoclonal antibody was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).


Secondary antibody only control: Secondary antibody is **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Immunohistochemical analysis of paraffin-embedded Human lung carcinoma tissue labeling Hsp70 with ab5439 at 1/2000 (0.519 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on human lung carcinoma. The section was incubated with ab5439 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.


Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized NIH/3T3 cells labelling Hsp70 with ab5439 at 1/50 (20.76 µg/ml) dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic and weak nuclear staining in NIH/3T3 cells **ab179513** Anti-beta Tubulin rabbit monoclonal antibody was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp70 antibody [3A3] (ab5439)

Immunohistochemical analysis of paraffin-embedded Mouse testis tissue labeling Hsp70 with ab5439 at 1/2000 (0.519 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on mouse testis. The section was incubated with ab5439 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).


Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins



Immunocytochemistry/ Immunofluorescence - Anti-Hsp70 antibody [3A3] (ab5439)

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized PC-12 cells labelling Hsp70 with ab5439 at 1/50 (20.76 µg/ml) dilution, followed by **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in PC-12 cells **ab179513** Anti-beta Tubulin rabbit monoclonal antibody was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150113** Goat Anti-Mouse IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.







Immunohistochemical analysis of paraffin-embedded Rat cerebrum tissue labeling Hsp70 with ab5439 at 1/2000 (0.519 µg/ml) dilution followed by a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection). Positive staining on rat cerebrum. The section was incubated with ab5439 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use LeicaDS9800 (Bond™ Polymer Refine Detection).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins

Why choose a recombinant antibody?

| | |
|--|--|
|  <p>Research with confidence Consistent and reproducible results</p> |  <p>Long-term and scalable supply Recombinant technology</p> |
|  <p>Success from the first experiment Confirmed specificity</p> |  <p>Ethical standards compliant Animal-free production</p> |

Anti-Hsp70 antibody [3A3] (ab5439)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors