

Anti-Hsp90 beta antibody [K3701] ab82584

敲除验证

3 图像

概述

产品名称	Anti-Hsp90 beta抗体[K3701]
描述	小鼠单克隆抗体[K3701] to Hsp90 beta
宿主	Mouse
经测试应用	适用于: Flow Cyt (Intra), WB
种属反应性	与反应: Rat, Human, Chinese hamster
免疫原	Recombinant human Hsp90 beta protein
阳性对照	WB: A431, Saos-2, HL-60, HeLa, PC12, CHO-K1 cell lysates; Hsp90 native protein. Flow Cyt (Intra): Human colon cancer Caco-2 cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	Preservative: 0.1% Sodium azide Constituent: PBS
纯度	Ammonium Sulphate Precipitation
克隆	单克隆
克隆编号	K3701
同种型	IgM

应用

The Abpromise guarantee


Abpromise™承诺保证使用ab82584于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use a concentration of 10 µg/ml. ab91545 - Mouse monoclonal IgM, is suitable for use as an isotype control with this antibody.
WB		1/1000. Predicted molecular weight: 83 kDa. 1/1000 (colorimetric).

靶标	
功能	Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function.
序列相似性	Belongs to the heat shock protein 90 family.
结构域	The TPR repeat-binding motif mediates interaction with TPR repeat-containing proteins.
翻译后修饰	Ubiquitinated in the presence of STUB1-UBE2D1 complex (in vitro). ISGylated. S-nitrosylated; negatively regulates the ATPase activity.
细胞定位	Cytoplasm. Melanosome. Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

图片



Western blot - Anti-Hsp90 beta antibody [K3701] (ab82584)

All lanes : Anti-Hsp90 beta antibody [K3701] (ab82584) at 1/1000 dilution

Lane 1 : Wild-type A-431 (Human epidermoid carcinoma cell line) whole cell lysate

Lane 2 : HSP90AB1 knockout A-431 (Human epidermoid carcinoma cell line) whole cell lysate

Lane 3 : Saos-2 (Human osteosarcoma cell line) whole cell lysate

Lane 4 : HL-60 (Human promyelocytic leukemia cell line) whole cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

Predicted band size: 83 kDa

Observed band size: 85 kDa

Lanes 1 - 4: Merged signal (red and green). Green - ab82584 observed at 85 kDa. Red - loading control, **ab181602** (Rabbit Anti-GAPDH antibody [EPR16891]) observed at 37kDa.

ab82584 was shown to react with Hsp90 beta in wild-type A431 cells in western blot. Loss of signal was observed when HSP90AB1 knockout sample was used. Wild-type A431 and HSP90AB1 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab82584 and **ab181602** (Rabbit Anti-GAPDH antibody [EPR16891]) overnight at 4°C at a 1 in 1000 Dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-Hsp90 beta antibody [K3701] (ab82584)

All lanes : Anti-Hsp90 beta antibody [K3701] (ab82584) at 1/1000 dilution

Lane 1 : HeLa cell lysate

Lane 2 : PC12 cell lysate


Lane 3 : CHO-K1 cell lysate

Lane 4 : Hsp90 Native Protein

Lane 5 : Hsp90 alpha Recombinant Protein (negative control)

Predicted band size: 83 kDa

Observed band size: 83 kDa



Flow Cytometry (Intracellular) - Anti-Hsp90 beta antibody [K3701] (ab82584)

ab82584, at 10 µg/ml, staining p90 beta in human colon cancer Caco-2 cells by flow cytometry (right). Isotype control antibody (left).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors