

# Anti-HTF9C/TRMT2A antibody ab121126

## 4 图像

### 概述

产品名称	Anti-HTF9C/TRMT2A抗体
描述	兔多克隆抗体to HTF9C/TRMT2A
宿主	Rabbit
经测试应用	适用于: ICC/IF, IHC-P, WB
种属反应性	与反应: Human
免疫原	Recombinant fragment corresponding to Human HTF9C/TRMT2A aa 462-621. Database link: <a href="#">Q8IZ69</a>
阳性对照	WB:RT4, EFO-21, HEK293 and A431 cell lysates; IHC-P: Human oral mucosa tissue; ICC/IF: U-251MG cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine)
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

### 应用

The Abpromise guarantee


Abpromise™承诺保证使用ab121126于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		Use a concentration of 0.25 - 2 µg/ml.
IHC-P		1/50 - 1/200. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
WB		Use a concentration of 0.04 - 0.4 µg/ml. Predicted molecular weight: 69 kDa.


靶标	
功能	May be involved in nucleic acid metabolism and/or modifications.
序列相似性	Contains 1 RRM (RNA recognition motif) domain.

图片




Anti-HTF9C/TRMT2A antibody (ab121126) at 0.4 µg/ml + HEK-293 cell lysate

**Predicted band size:** 69 kDa




Immunohistochemical analysis of paraffin-embedded human oral mucosa staining HTF9C/TRMT2A with ab121126 at 1/50 dilution.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-HTF9C/TRMT2A antibody (ab121126)



Immunofluorescent staining of Human cell line U-251MG shows positivity in nucleus and cytoplasm. Recommended concentration of ab121126 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-HTF9C/TRMT2A antibody (ab121126)



**All lanes :** Anti-HTF9C/TRMT2A antibody (ab121126) at 1/250 dilution

**Lane 1 :** RT4 cell lysate

**Lane 2 :** EFO-21 cell lysate

**Lane 3 :** A431 cell lysate

**Lane 4 :** Human Liver lysate

**Lane 5 :** Human Tonsil lysate

Developed using the ECL technique.

**Predicted band size:** 69 kDa

10-20% SDS-PAGE.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors