# abcam

# Product datasheet

# Anti-IKB alpha (phospho Y305) antibody ab24784

★★★☆☆ 1 Abreviews 2 References 1 图像

概述

产品名称 Anti-IKB alpha (phospho Y305)抗体

描述 兔多克隆抗体to IKB alpha (phospho Y305)

**宿主** Rabbit

经测试应用 适用于: WB

种属反应性 与反应: Human

免疫原 Synthetic peptide corresponding to Human IKB alpha (phospho Y305) conjugated to Keyhole

Limpet Haemocyanin (KLH). Synthetic peptide conjugated to KLH corresponding to amino acid residues around tyrosine 305 of human IKB alpha. This peptide sequence has low homology to

other IKB proteins.

阳性对照 A431 and Jurkat cells treated with pervanadate.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or

contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Liquid

**存放说明** Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

存储溶液 Preservative: 0.05% Sodium azide

Constituents: PBS, 50% Glycerol, 0.1% BSA

纯**度** Immunogen affinity purified

纯**化说明** This antibody was cross-adsorbed to phospho-tyrosine coupled to agarose then affinity purified

using phospho IKB alpha (Tyr-305) peptide.

**克隆** 多克隆

同种型 lgG

1

## The Abpromise guarantee

## Abpromise™承诺保证使用ab24784于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB	<b>★★★</b> ☆☆ (1)	1/500. Detects a band of approximately 38 kDa.

#### 靶标

## 功能

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.

# 疾病相关

# 序列相似性

Ectodermal dysplasia, anhidrotic, with T-cell immunodeficiency autosomal dominant

Belongs to the NF-kappa-B inhibitor family.

Contains 5 ANK repeats.

## 翻译后修饰

Phosphorylated; disables inhibition of NF-kappa-B DNA-binding activity. Phosphorylation at positions 32 and 36 is prerequisite to recognition by UBE2D3 leading to polyubiquitination and subsequent degradation.

Sumoylated; sumoylation requires the presence of the nuclear import signal. Sumoylation blocks ubiquitination and proteasome-mediated degradation of the protein thereby increasing the protein stability.

Monoubiquitinated at Lys-21 and/or Lys-22 by UBE2D3. Ubiquitin chain elongation is then performed by CDC34 in cooperation with the SCF(FBXW11) E3 ligase complex, building ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. The resulting polyubiquitination leads to protein degradation. Also ubiquitinated by SCF(BTRC) following stimulus-dependent phosphorylation at Ser-32 and Ser-36.


Deubiquitinated by porcine reproductive and respiratory syndrome virus Nsp2 protein, which thereby interferes with NFKBIA degradation and impairs subsequent NF-kappa-B activation.

#### 细胞定位

Cytoplasm. Nucleus. Shuttles between the nucleus and the cytoplasm by a nuclear localization

signal (NLS) and a CRM1-dependent nuclear export.

### 图片



Western blot - Anti-IKB alpha (phospho Y305) antibody (ab24784)

Western blot analysis of A431 cells treated with pervanadate (1 mM) for 30 minutes (30 ug/lane). Blots were probed with antilkappaBalpha (lane 1), anti-phospho-lkappaBalpha?(Tyr-42) (ab24783; lanes 2-5), or anti-phospho-lkappaBalpha (Tyr-305) (ab24784; lanes 6-9). Both anti-phospho-lkappaBalpha (Tyr-42) and anti-phospho-lkappaBalpha (Tyr-305) were used in the presence of no blocking peptide (lane 2 & 6), phospho-lkappaBalpha (Tyr-42) peptide (lane 3 & 8), phospho-lkappaBalpha (Tyr-305) peptide (lane 4 & 7), or BSA conjugated to phosphotyrosine (lane 5 & 9). Peptides and BSA-pTyr were used at 1 ug/ml.

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

## Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors