# abcam

## Product datasheet

## Anti-Integrin linked ILK antibody [EP1593Y] ab52480





重组 RabMAb

18 References 10 图像

概述

产品名称 Anti-Integrin linked LK抗体[EP1593Y]

描述 兔单克隆抗体[EP1593Y] to Integrin linked ILK

宿主 Rabbit

特异性 This antibody reacts with Integrin linked ILK. The mouse and rat recommendation is based on the

WB results. We do not guarantee IHC-P for mouse and rat.

适用于: WB, IP, IHC-P 经测试应用

不适用于: Flow Cyt or ICC/IF

种属反应性 与反应: Rat, Human, African green monkey

预测可用于: Mouse, Cow 📤

免疫原 Synthetic peptide within Human Integrin linked ILK aa 400 to the C-terminus. The exact sequence

is proprietary.

Database link: Q13418

阳性对照 Human heart muscle tissue and COS cell lysate

常规说明 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

性能

形式 Liquid

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

存储溶液 pH: 7.20

Preservative: 0.01% Sodium azide

Constituents: 59% PBS, 40% Glycerol, 0.05% BSA

纯**度** Protein A purified

**克隆** 单克隆

**克隆编号** EP1593Y

同种型 lgG

#### 应用

### The Abpromise guarantee Abpromise™承诺保证使用ab52480于以下的经测试应用

"应用说明"部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Detects a band of approximately 51 kDa (predicted molecular weight: 51 kDa).  For unpurified use at 1/2000 - 1/10000 dilution.
IP		1/40. For unpurified use at 1/30 dilution.
IHC-P		1/100. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.  See IHC antigen retrieval protocols.  The mouse and rat recommendation is based on the WB results.  We do not guarantee IHC-P for mouse and rat.

应用说明 Is unsuitable for Flow Cyt or ICC/IF.

#### 靶标

序列相似性 Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.

Contains 5 ANK repeats.


Contains 1 protein kinase domain.

结**构域** A PH-like domain is involved in phosphatidylinositol phosphate binding.


翻译后修饰 Autophosphorylated on serine residues.

细**胞定位** Cell junction > focal adhesion. Cell membrane.

## 图片



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: Integrin linked ILK knockout HAP1 cell lysate (20 µg)

Lane 3: HEK293 cell lysate (20 µg)

Lane 4: K562 cell lysate (20 µg)

**Lanes 1 - 4:** Merged signal (red and green). Green - ab52480 observed at 52 kDa. Red - loading control, <u>ab8245</u>, observed at 37 kDa.

Unpurified ab52480 was shown to specifically react with Integrin linked ILK when Integrin linked ILK knockout samples were used. Wild-type and Integrin linked ILK knockout samples were subjected to SDS-PAGE. Unpurified ab52480 and <a href="mailto:ab8245">ab8245</a> (loading control to GAPDH) were both diluted 1/2000 and incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (<a href="mailto:ab216773">ab216773</a>) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (<a href="mailto:ab216776">ab216776</a>) secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human lung cancer tissue sections labeling Integrin linked ILK with Purified ab52480 at 1/100 dilution (7.13 µg/ml). Heat mediated antigen retrieval was performed using citrate (pH 6.0).



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

**All lanes :** Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/1000 dilution (Purified)

**Lane 1**: HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysates with 5% NFDM/TBST

**Lane 2 :** K-562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysates with 5% NFDM/TBST

Lane 3: Mouse heart lysates with 5% NFDM/TBST

**Lane 4**: HEK-293 (Human embryonic kidney epithelial cell) whole cell lysates with 5% NFDM/TBST

**Lane 5**: PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysates with 5% NFDM/TBST

Lysates/proteins at 20 µg per lane.

## **Secondary**

**All lanes :** Goat Anti-Rabbit lgG H&L (HRP) (<u>ab97051</u>) at 1/20000 dilution (Goat Anti-Rabbit lgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 51 kDa **Observed band size:** 51 kDa



Immunoprecipitation - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

ab52480 (purified) at 1:40 dilution (2µg) immunoprecipitating Integrin linked ILK in HeLa whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) whole cell lysate 10µg


Lane 2 (+): ab52480 & HeLa whole cell lysate

**Lane 3 (-):** Rabbit monoclonal IgG (<u>ab172730</u>) instead of ab52480 in HeLa whole cell lysate

For western blotting, VeriBlot for IP Detection Reagent (HRP)

(ab131366) was used for detection at 1:1000 dilution.

Blocking and diluting buffer: 5% NFDM/TBST.




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/1000 dilution (Purified) + COS-1 (African green monkey kidney fibroblast-like) whole cell lysates at 15 µg with 5% NFDM/TBST

## **Secondary**

Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/20000 dilution (Goat Anti-Rabbit lgG, (H+L), Peroxidase conjugated)

**Predicted band size:** 51 kDa **Observed band size:** 51 kDa




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) at 1/2000 dilution (unpurified) + COS-1 (African green monkey kidney fibroblast-like) cell lysate at 10 µg

## Secondary

Goat anti-rabbit HRP labeled at 1/2000 dilution


**Predicted band size:** 51 kDa **Observed band size:** 51 kDa



Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)


Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) (unpurified) + PC12

Predicted band size: 51 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

Immunohistochemical analysis of paraffin-embedded human heart muscle using unpurified ab52480 at a 1/50 dilution.




Western blot - Anti-Integrin linked ILK antibody [EP1593Y] (ab52480)

 $\begin{tabular}{ll} \textbf{All lanes:} Anti-Integrin linked ILK antibody [EP1593Y] (ab52480) \\ at 1/2000 dilution (unpurified) \\ \end{tabular}$ 

Lane 1: Rat Brain
Lane 2: Rat Heart
Lane 3: Rat Kidney
Lane 4: Rat Spleen

Predicted band size: 51 kDa

Exposure time: 3 minutes



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- · We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.cn/abpromise">https://www.abcam.cn/abpromise</a> or contact our technical team.

### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors