


### Anti-Islet 1 antibody ab20670

★★★★★ [17 Abreviews](#) [83 References](#) [9 图像](#)

#### 概述

产品名称	Anti-Islet 1抗体
描述	兔多克隆抗体to Islet 1
宿主	Rabbit
特异性	<p>ab20670 might also detect rat and human Islet 2 protein as the immunogen used to raise this ab20670 is 86% identical to rat and human Islet 2. This has not been tested.</p> <p>Replenishment batches of our polyclonal antibody, ab20670 are tested in IHC-P. Previous batches were additionally validated in ICC/IF and IHC-Fr. These applications are still expected to work and are covered by our Abpromise guarantee. You may also be interested in our alternative recombinant antibody, <a href="#">ab109517</a>.</p>
经测试应用	<b>适用于:</b> IHC-P, ICC/IF, IHC-Fr
种属反应性	<p><b>与反应:</b> Mouse, Rat, Human, Apteronotus leptorhynchus</p> <p><b>预测可用于:</b> Chicken, Zebrafish </p>
免疫原	<p>Synthetic peptide corresponding to Human Islet 1 aa 300 to the C-terminus (C terminal) conjugated to keyhole limpet haemocyanin.</p> <p>(Peptide available as <a href="#">ab21995</a>, <a href="#">ab21996</a>)</p>
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

#### 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	<p>pH: 7.40</p> <p>Preservative: 0.02% Sodium azide</p> <p>Constituent: PBS</p>

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

## 应用

**The Abpromise guarantee**      **Abpromise™**承诺保证使用ab20670于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (3)	Use a concentration of 0.5 - 1 µg/ml. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF	★★★★★ (3)	Use a concentration of 2 µg/ml.
IHC-Fr	★★★★★ (5)	1/500.

## 靶标

功能	Binds to one of the cis-acting domain of the insulin gene enhancer.
组织特异性	Expressed in subsets of neurons of the adrenal medulla and dorsal root ganglion, inner nuclear and ganglion cell layers in the retina, the pineal and some regions of the brain.
序列相似性	Contains 1 homeobox DNA-binding domain. Contains 2 LIM zinc-binding domains.
细胞定位	Nucleus.


## 图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Islet 1 antibody (ab20670)

IHC image of Islet 1 staining in a section of formalin fixed, paraffin embedded mouse embryo E12, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval (EDTA based pH 9.0 solution, epitope retrieval solution 2) for 20 mins. The section was then incubated with ab20670, 0.5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Islet 1 antibody (ab20670)

IHC image of Islet 1 staining in a section of formalin fixed, paraffin embedded mouse embryo E12, performed on a Leica Bond™ system using the standard protocol B. The section was pre-treated using heat mediated antigen retrieval (EDTA based pH 9.0 solution, epitope retrieval solution 2) for 20 mins. The section was then incubated with ab20670, 0.5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.


For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Islet 1 antibody (ab20670)

ab20670 2µg/ml staining ISLET1 in human pancreas using an automated system (DAKO Autostainer Plus). Using this protocol there is strong nuclear and weak cytoplasmic staining primarily in the pancreatic islet.

Sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffer EDTA pH 9.0 in a DAKO PT Link. Slides were peroxidase blocked in 3% H<sub>2</sub>O<sub>2</sub> in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako Envision Flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX. Please note that, for manual staining, optimization of primary antibody concentration and incubation time is recommended. Signal amplification may be required.



Immunocytochemistry/ Immunofluorescence - Anti-Islet 1 antibody (ab20670)

This image is courtesy of Randal Moldrich, CNRS UMR7637, ESPCI, France


Dorsal root ganglion explants were dissected from 16 day-old rat embryos and cultured for 6 hours in vitro with Neurobasal Medium containing B27 supplement.

Nuclei stained positive for anti-Islet 1 antibody ab20670 at 2µg/ml. As would be expected, not all cells in this preparation were Islet 1-positive. Pre-incubation of ab20670 with the immunizing peptide **ab21996** resulted in complete blocking of the antibody.

Green = ab20670

Blue = To-pro-3 nuclear stain


The level of magnification is different in each image.



Immunocytochemistry/ Immunofluorescence - Anti-Islet 1 antibody (ab20670)

Image from Ifkovits, Jamie L. et al. PLoS ONE 9.2 (2014): e89678. doi: 10.1371/journal.pone.0089678. Fig S3B. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>


Immunocytochemistry/ Immunofluorescence analysis of adult mouse cardiac fibroblasts labeling Islet 1 with ab20670 at 1/100 dilution (green). Samples were fixed using 4% PFA with 0.25% TritonX-100. Scale bar is 100  $\mu$ M.



Immunohistochemistry (Frozen sections) - Anti-Islet 1 antibody (ab20670)

This image is courtesy of an anonymous abreview.

Immunohistochemistry (Frozen sections) analysis of mouse brain tissue sections labeling Islet 1 with ab20670 at 1/200 dilution. The tissue was fixed with paraformaldehyde followed by blocking with 5% serum for 1 hour at 25°C. The tissue was incubated with ab20670 in PBST for 12 hours at 4°C. A polyclonal donkey anti-rabbit Alexa Fluor® 594 secondary antibody was used at 1/1000 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-Islet 1 antibody (ab20670)

This image is courtesy of an anonymous abreview.

Immunocytochemistry/ Immunofluorescence analysis of mouse dorsal root ganglia (DRG) neurons labeling Islet 1 with ab20670 at 1/200 dilution. Cells were fixed with paraformaldehyde and permeabilized with Tween 20. Blocking of the cells was performed with 2% serum for 30 minutes at 21°C, followed by incubation with ab20670 for 18 hours at 4°C. A polyclonal goat anti-rabbit Alexa Fluor® 568 secondary antibody was used at 1/1000 dilution. DAPI was used to counterstain.






Immunohistochemical analysis of paraffin-embedded mouse E11.5 somites labelling Islet 1 with ab20670 at 1/100 dilution, followed by BioGenex polymer-HRP reagent (undiluted according to manufacturer).

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Islet 1 antibody (ab20670)

This image is courtesy of an anonymous collaborator.



ab20670 detected Islet 1 in the nucleus of frozen sections of E11.5 mouse embryonic brain using 1 ug/ml of antibody. Brighter staining may be achieved using a greater concentration of antibody, e.g. 2.5 or 5 ug/ml.

Immunohistochemistry (Frozen sections) - Anti-Islet 1 antibody (ab20670)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors