

Anti-ITGA7 antibody [3C12] ab195959

[2 References](#) [2 图像](#)

概述

产品名称	Anti-ITGA7抗体[3C12]
描述	小鼠单克隆抗体[3C12] to ITGA7
宿主	Mouse
经测试应用	适用于: Flow Cyt, IHC-P
种属反应性	与反应: Mouse, Human
免疫原	Tissue, cells or virus corresponding to Mouse ITGA7. ab195959 was raised against Mouse myoblasts.
阳性对照	Human brain, cortex tissue; C2C12 cells.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
存储溶液	Constituents: 50% Glycerol (glycerin, glycerine), 50% PBS
纯度	Protein A purified
克隆	单克隆
克隆编号	3C12
同种型	IgG1

应用

The Abpromise guarantee

Abpromise™ 承诺保证使用ab195959于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt		Use a concentration of 10 - 20 µg/ml. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
IHC-P		Use a concentration of 10 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标


功能	<p>Integrin alpha-7/beta-1 is the primary laminin receptor on skeletal myoblasts and adult myofibers. During myogenic differentiation, it may induce changes in the shape and mobility of myoblasts, and facilitate their localization at laminin-rich sites of secondary fiber formation. It is involved in the maintenance of the myofibers cytoarchitecture as well as for their anchorage, viability and functional integrity. Isoform Alpha-7X2B and isoform Alpha-7X1B promote myoblast migration on laminin 1 and laminin 2/4, but isoform Alpha-7X1B is less active on laminin 1 (In vitro). Acts as Schwann cell receptor for laminin-2. Acts as a receptor of COMP and mediates its effect on vascular smooth muscle cells (VSMCs) maturation (By similarity). Required to promote contractile phenotype acquisition in differentiated airway smooth muscle (ASM) cells.</p>
组织特异性	<p>Isoforms containing segment A are predominantly expressed in skeletal muscle. Isoforms containing segment B are abundantly expressed in skeletal muscle, moderately in cardiac muscle, small intestine, colon, ovary and prostate and weakly in lung and testes. Isoforms containing segment X2D are expressed at low levels in fetal and adult skeletal muscle and in cardiac muscle, but are not detected in myoblasts and myotubes. In muscle fibers isoforms containing segment A and B are expressed at myotendinous and neuromuscular junctions; isoforms containing segment C are expressed at neuromuscular junctions and at extrasynaptic sites. Isoforms containing segments X1 or X2 or, at low levels, X1X2 are expressed in fetal and adult skeletal muscle (myoblasts and myotubes) and cardiac muscle.</p>
疾病相关	<p>Defects in ITGA7 are the cause of muscular dystrophy congenital due to integrin alpha-7 deficiency (MDCI) [MIM:613204]. A form of congenital muscular dystrophy. Patients present at birth, or within the first few months of life, with hypotonia, muscle weakness and often with joint contractures.</p>
序列相似性	<p>Belongs to the integrin alpha chain family. Contains 7 FG-GAP repeats.</p>
发展阶段	<p>In renewing intestinal epithelium, expression of isoforms containing segment B correlates with the onset of enterocytic differentiation.</p>
翻译后修饰	<p>ADP-ribosylated on at least two sites of the extracellular domain in skeletal myotubes. A 70 kDa form is created by proteolytic cleavage. Cleavage is elevated during myogenic differentiation and the cleaved form enhances cell adhesion and spreading on laminin.</p>
细胞定位	<p>Membrane.</p>

图片



Flow Cytometry - Anti-ITGA7 antibody [3C12]
(ab195959)

Flow Cytometrical analysis of C2C12 cells labeling ITGA7 with ab195959 at 20 μg or an isotypic control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ITGA7 antibody [3C12]
(ab195959)

Immunohistochemical analysis of formalin-fixed, paraffin-embedded human brain, cortex tissue labeling ITGA7 with ab195959 at 10 $\mu\text{g}/\text{mL}$.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors