

Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] ab32101

重组 RabMAb

★★★★☆ **4 Abreviews** **171 References** [12 图像](#)

概述

产品名称	Anti-JAK2 (phospho Y1007 + Y1008)抗体[E132]
描述	兔单克隆抗体[E132] to JAK2 (phospho Y1007 + Y1008)
宿主	Rabbit
特异性	<p>This antibody is phospho-specific and only detects phosphorylated JAK2 on Tyrosine 1007 and 1008 (pY1007+Y1008). According to our ELISA results, this antibody preferentially recognizes phospho Y1007. Stimulation may be required to allow detection of the phosphorylated protein. Please see images below for recommended treatment conditions and positive controls.</p> <p>The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.</p>
经测试应用	适用于: ELISA, Flow Cyt (Intra), WB, ICC/IF, IHC-P, Dot blot
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: Hepa1-6, MEF, C6, Jurkat treated with Pervanadate and Jurkat cell lysates. IHC-P: Human differentiated squamous cell carcinoma tissue. ICC/IF: Jurkat cells (treated with Pervanadate). Flow Cyt (intra): Jurkat starved of serum for 16 hours then treated with 1mM Pervanadate for 30 minutes.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.

存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	E132
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab32101于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
Flow Cyt (Intra)		1/20.
WB	★★★★★ (3)	1/1000 - 1/10000. Detects a band of approximately 120 kDa (predicted molecular weight: 130 kDa). The samples may require stimulation (E.g., Jurkat cells treated with pervanadate for 5 min)
ICC/IF	★★★★★ (1)	1/1000.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat.
Dot blot		1/1000.

靶标

功能

Non-receptor tyrosine kinase involved in various processes such as cell cycle progression, apoptosis, mitotic recombination, genetic instability and histone modifications. In the cytoplasm, plays a pivotal role in signal transduction via its association with cytokine receptors, which constitutes an initiating step in signaling for many members of the cytokine receptor superfamily including the receptors for growth hormone (GHR), prolactin (PRLR), leptin (LEPR), erythropoietin (EPOR), granulocyte-macrophage colony-stimulating factor (CSF2), thrombopoietin (THPO) and multiple interleukins. Following stimulation with erythropoietin (EPO) during erythropoiesis, it is autophosphorylated and activated, leading to its association with erythropoietin receptor (EPOR) and tyrosine phosphorylation of residues in the EPOR cytoplasmic domain. Also involved in promoting the localization of EPOR to the plasma membrane. Also acts downstream of some G-protein coupled receptors. Plays a role in the control of body weight (By similarity). Mediates angiotensin-2-induced ARHGEF1 phosphorylation. In the nucleus, plays a key role in chromatin by specifically mediating phosphorylation of 'Tyr-41' of histone H3 (H3Y41ph), a specific tag that

promotes exclusion of CBX5 (HP1 alpha) from chromatin.

组织特异性

Expressed in blood, bone marrow and lymph node.

疾病相关

Note=Chromosomal aberrations involving JAK2 are found in both chronic and acute forms of eosinophilic, lymphoblastic and myeloid leukemia. Translocation t(8;9)(p22;p24) with PCM1 links the protein kinase domain of JAK2 to the major portion of PCM1. Translocation t(9;12)(p24;p13) with ETV6.

Defects in JAK2 are a cause of susceptibility to Budd-Chiari syndrome (BCS) [MIM:600880]. It is a syndrome caused by obstruction of hepatic venous outflow involving either the hepatic veins or the terminal segment of the inferior vena cava. Obstructions are generally caused by thrombosis and lead to hepatic congestion and ischemic necrosis. Clinical manifestations observed in the majority of patients include hepatomegaly, right upper quadrant pain and abdominal ascites. Budd-Chiari syndrome is associated with a combination of disease states including primary myeloproliferative syndromes and thrombophilia due to factor V Leiden, protein C deficiency and antithrombin III deficiency. Budd-Chiari syndrome is a rare but typical complication in patients with polycythemia vera.

Defects in JAK2 are a cause of polycythemia vera (PV) [MIM:263300]. A myeloproliferative disorder characterized by abnormal proliferation of all hematopoietic bone marrow elements, erythroid hyperplasia, an absolute increase in total blood volume, but also by myeloid leukocytosis, thrombocytosis and splenomegaly.

Defects in JAK2 gene may be a cause of essential thrombocythemia (ET) [MIM:187950]. ET is characterized by elevated platelet levels due to sustained proliferation of megakaryocytes, and frequently lead to thrombotic and haemorrhagic complications.

Defects in JAK2 are a cause of myelofibrosis (MYELOF) [MIM:254450]. Myelofibrosis is a disorder characterized by replacement of the bone marrow by fibrous tissue, occurring in association with a myeloproliferative disorder. Clinical manifestations may include anemia, pallor, splenomegaly, hypermetabolic state, petechiae, ecchymosis, bleeding, lymphadenopathy, hepatomegaly, portal hypertension.

Defects in JAK2 are a cause of acute myelogenous leukemia (AML) [MIM:601626]. AML is a malignant disease in which hematopoietic precursors are arrested in an early stage of development.

序列相似性

Belongs to the protein kinase superfamily. Tyr protein kinase family. JAK subfamily.

Contains 1 FERM domain.

Contains 1 protein kinase domain.

Contains 1 SH2 domain.

结构域

Possesses 2 protein kinase domains. The second one probably contains the catalytic domain, while the presence of slight differences suggest a different role for protein kinase 1.


翻译后修饰

Autophosphorylated, leading to regulate its activity. Leptin promotes phosphorylation on tyrosine residues, including phosphorylation on Tyr-813. Autophosphorylation on Tyr-119 in response to EPO down-regulates its kinase activity. Autophosphorylation on Tyr-868, Tyr-966 and Tyr-972 in response to growth hormone (GH) are required for maximal kinase activity.

细胞定位

Endomembrane system. Nucleus.


图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

ab32101 showing positive staining in Human differentiated squamous cell carcinoma of the cervix tissue at 1/10000 dilution. Goat Anti-Rabbit IgG H&L (HRP) was used as secondary antibody. Antigen retrieval was carried out by Heat mediated antigen retrieval using **ab93684** (Tris/EDTA buffer, pH 9.0).

Nuclear staining on human differentiated squamous cell carcinoma of the cervix without alkaline phosphatase treatment (image A). No staining on human differentiated squamous cell carcinoma of the cervix with alkaline phosphatase treatment (image B)



Western blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

All lanes : Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101) at 1/1000 dilution

Lane 1 : Hepa1-6 (Mouse hepatoma epithelial cell) whole cell lysate

Lane 2 : Hepa1-6 (Mouse hepatoma epithelial cell) treated with 100 μM pervanadate for 30 minutes whole cell lysate

Lane 3 : MEF (Mouse embryonic fibroblast (immortalized)) whole cell lysate

Lane 4 : MEF (Mouse embryonic fibroblast (immortalized)) treated with 100 μM pervanadate for 30 minutes whole cell lysate


Lysates/proteins at 15 μg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 130 kDa

The extra bands are undefined.



Western blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

All lanes : Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)


- Lane 1** : Mouse hippocampus lysate
- Lane 2** : Mouse P240 hippocampus lysate
- Lane 3** : Mouse P7 hippocampus lysate
- Lane 4** : Rat hippocampus lysate
- Lane 5** : Rat P7 hippocampus lysate
- Lane 6** : Rat brain cortex lysate
- Lane 7** : Human brain lysate
- Lane 8** : Mouse brain lysate
- Lane 9** : Rat brain lysate
- Lane 10** : C6 (Rat glial tumor glial cell) whole cell lysate
- Lane 11** : C6 (Rat glial tumor glial cell) treated with 50mM pervanadate for 5 minutes whole cell lysate

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution (Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated)

Predicted band size: 130 kDa



Western blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

All lanes : Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101) at 1/5000 dilution

- Lane 1** : Untreated Jurkat cells whole cell lysates
- Lane 2** : Jurkat cells were treated with 50mM Pervanadate for 5 minutes whole cell lysates
- Lane 3** : Jurkat cells were treated with 50mM Pervanadate for 5 minutes whole cell lysates. Then the membrane was incubated with Alkaline phosphatase.

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 130 kDa

Observed band size: 120 kDa


Exposure time: 5 seconds

Blocking and diluting buffer 5% NFDN/TBST



Immunocytochemistry/ Immunofluorescence - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)


Immunocytochemistry/Immunofluorescence analysis of Jurkat +/- pervanadate (1mM, 30min) and Jurkat + pervanadate (1mM, 30min) + LP cells labelling JAK2 (phospho Y1007 + Y1008) with ab32101 at a dilution of 1/1000. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% tritonX-100. **ab150077** (goat anti-rabbit IgG Alexa Fluor[®] 488) (1/1000) was used as the secondary antibody. The cells were co-stained with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594)) at a 1/200 dilution. Nuclei counterstained with DAPI (blue).



Dot Blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

Dot blot analysis of human JAK2 (phospho Y1007 & Y1008) phospho peptide (Lane 1), JAK2 (phospho Y1007) phospho peptide (Lane 2), JAK2 (phospho Y1008) phospho peptide (Lane 3) and JAK2 non-phospho peptide (Lane 4) labelling JAK2 (phospho Y1007 & Y1008) with ab32101 at a dilution of 1/1000. Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (**ab97051**) was used as the secondary antibody at a dilution of 1/20,000.

Blocking/Dilution buffer: 5% NFDN/TBST.



Western blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

All lanes : Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101) at 1/1000 dilution

Lane 1 : Jurkat (Human T cell leukemia T lymphocyte) whole cell lysates

Lane 2 : Jurkat (Human T cell leukemia T lymphocyte) treated with 50mM Pervanadate for 5 minutes whole cell lysates

Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) (**ab97051**) at 1/20000 dilution

Predicted band size: 130 kDa

Observed band size: 120 kDa


Additional bands at: 60 kDa. We are unsure as to the identity of these extra bands.

Blocking and diluting buffer: 5% NFDM/TBST

Exposure time:

Left image: 1 second


Right image: 5 minutes



Flow Cytometry (Intracellular) - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

Intracellular Flow Cytometry analysis of Jurkat (human acute T cell leukemia) cells starved of serum for 16 hours then treated with 1 mM Pervanadate for 30 minutes labeling JAK2 (phospho Y1007 + Y1008) with ab32101 at 1/20 dilution (10 µg/mL) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor®488) at 1/2000 dilution was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) was used as the unlabeled control. Unstimulated Jurkat cells were used as a negative control (Green).

Direct ELISA antibody dose-response curve at 100 ng/ml peptide




ELISA - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

Direct ELISA antigen dose-response curve using ab32101 at 0~1000 ng/mL. Antigen concentration of 100 ng/mL. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (1/2500) was used as the secondary antibody.

This antibody preferentially recognizes phospho Y1007. When the concentration of peptides is higher than 100 ng/mL, it also recognizes phospho Y1008.


Direct ELISA antibody dose-response curve at 10 ng/ml peptide



ELISA - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

Direct ELISA antigen dose-response curve using ab32101 at 0~1000 ng/mL. Antigen concentration of 10 ng/mL. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (1/2500) was used as the secondary antibody.

This antibody preferentially recognizes phospho Y1007. When the concentration of peptides is lower than 10 ng/mL, it cannot recognize phospho Y1008.



Western blot - Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101)

This image is courtesy of an anonymous Abreview

All lanes : Anti-JAK2 (phospho Y1007 + Y1008) antibody [E132] (ab32101) at 1/2000 dilution

Lane 1 : Rat uterine cell line - whole cell lysate. Treated with 1 µg/mL Prolactin for 0 hours.

Lane 2 : Rat uterine cell line - whole cell lysate. Treated with 1 µg/mL Prolactin for 15 minutes.

Lane 3 : Rat uterine cell line - whole cell lysate. Treated with 1 µg/mL Prolactin for 30 minutes.

Lane 4 : Rat uterine cell line - whole cell lysate. Treated with 1 µg/mL Prolactin for 1 hour.

Lane 5 : Rat uterine cell line - whole cell lysate. Treated with 1µg/mL Prolactin for 2 hours.

Lane 6 : Rat uterine cell line - whole cell lysate. Treated with 1µg/mL Prolactin for 4 hours.

Lane 7 : Rat uterine cell line - whole cell lysate. Treated with 1µg/mL Prolactin for 6 hours.

Lane 8 : Rat uterine cell line - whole cell lysate. Treated with 1µg/mL Prolactin for 24 hours.

Lysates/proteins at 30 µg per lane.

Secondary

All lanes : An HRP-conjugated donkey anti-rabbit polyclonal. at 1/10000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 130 kDa

Observed band size: 110 kDa

Additional bands at: 55 kDa (possible non-specific binding)

Why choose a recombinant antibody?



- Research with confidence**
Consistent and reproducible results
- Long-term and scalable supply**
Recombinant technology
- Success from the first experiment**
Confirmed specificity
- Ethical standards compliant**
Animal-free production

Anti-JAK2 (phospho Y1007 + Y1008) antibody
[E132] (ab32101)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Promise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet

- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors