


Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free ab219584

重组 RabMAb

13 References [7 图像](#)

概述

| | |
|-------|---|
| 产品名称 | Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221)抗体[EPR5693] - BSA and Azide free |
| 描述 | 兔单克隆抗体[EPR5693] to JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) - BSA and Azide free |
| 宿主 | Rabbit |
| 特异性 | This antibody will detect will detect JNK1 (pT183), JNK2 (pT183) and JNK3 (pT221). |
| 经测试应用 | 适用于: Flow Cyt (Intra), WB, IP, IHC-P, ICC/IF, Dot blot |
| 种属反应性 | 与反应: Mouse, Human 预测可用于: Rat  |
| 免疫原 | Synthetic peptide. This information is proprietary to Abcam and/or its suppliers. |
| 阳性对照 | NIH 3T3 cell lysates treated with Anisomycin; Human brain tissue. IP: HeLa treated with 25ug/mL anisomycin for 30min whole cell lysate. |
| 常规说明 | ab219584 is the carrier-free version of ab124956 . |

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:


- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb[®] patents](#).

性能

| | |
|------------------------|---|
| 形式 | Liquid |
| 存放说明 | Shipped at 4°C. Store at +4°C. Do Not Freeze. |
| 解离常数 (K _D) | K _D = 2.09 x 10 ⁻¹¹ M |



| | |
|------|-----------------------------|
| 存储溶液 | pH: 7.2 Constituent: PBS |
| 无载体 | 是 |
| 纯度 | Protein A purified |
| 克隆 | 单克隆 |
| 克隆编号 | EPR5693 |
| 同种型 | IgG |

应用

The Abpromise guarantee [Abpromise[™]](#) 承诺保证使用 ab219584 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度; 实际最佳的稀释度/浓度应由使用者检定。


| 应用 | Ab 评论 | 说明 |
|------------------|-------|--|
| Flow Cyt (Intra) | | Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |
| WB | | Use at an assay dependent concentration. Detects a band of approximately 46-54 kDa. |
| IP | | Use at an assay dependent concentration. |
| IHC-P | | Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol. (Heat to 98°C, allow to cool for 10-20 minutes) |
| ICC/IF | | Use at an assay dependent concentration. |
| Dot blot | | Use at an assay dependent concentration. |

靶标

细胞定位

Cytoplasmic, Mitochondrial, Nuclear and Plasma membrane

图片




Immunocytochemistry/Immunofluorescence analysis of untreated, Anisomycin treated and Anisomycin + LP treated NIH/3T3 cells labelling JNK1 + JNK2 + JNK3 (phospho T183 + T183 + T221) with **ab124956** at a dilution of 1/100 (left) and JNK1 + JNK2 + JNK3 with **ab179461** at a dilution of 1/250 (right).

Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/1000) were also used.

The image shows increased nuclear staining after Anisomycin (250ng/ml, 30min) treatment on NIH3T3 cells. The LP treatment decreased the increased nuclear staining caused by Anisomycin.

ab179461 was used as a Pan control for **ab124956**. The results showed cytoplasmic staining on untreated, Anisomycin and Anisomycin + LP treated NIH3T3 cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124956**).




Dot blot analysis of JNK1/2/3 (pT183 + pT183 + pT221) peptide (Lane 1) and JNK1/2/3 non-phospho peptide (Lane 2) labelling JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) with **ab124956** at a dilution of 1/1000. **ab97051** (Peroxidase conjugated goat anti-rabbit IgG (H+L)) was used as the secondary antibody at a dilution of 1/100000.

Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.


This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124956**).



Flow Cytometry (Intracellular) - Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free (ab219584)

Overlay histogram showing HeLa cells stained with **ab124956** (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (**ab124956**, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was goat anti-rabbit Alexa Fluor® 488 IgG (H+L) (**ab150077**) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124956**).



Immunoprecipitation - Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free (ab219584)

This data was developed using **ab124956**, the same antibody clone in a different buffer formulation.

Purified **ab124956** at 1/70 dilution (2µg) immunoprecipitating JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) in HeLa treated with 25ug/mL anisomycin for 30min whole cell lysate.

Lane 1 (input): HeLa (Human cervix adenocarcinoma epithelial cell) treated with 25ug/mL anisomycin for 30min whole cell lysate 10µg

Lane 2 (+): **ab124956** + HeLa treated with 25ug/mL anisomycin for 30min whole cell lysate.


Lane 3 (-): Rabbit monoclonal IgG (**ab172730**) instead of **ab124956** in HeLa treated with 25ug/mL anisomycin for 30min whole cell lysate.

VeriBlot for IP Detection Reagent (HRP) (**ab131366**) (1/5000 dilution) was used for Western blotting.

Blocking Buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.

Observed band size: 46, 54 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free (ab219584)

ab124956, at 1/100 dilution staining JNK1+JNK2+JNK3 in paraffin-embedded Human brain tissue, by Immunohistochemistry.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124956**).

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



OIR-D Scanning - Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free (ab219584)

Equilibrium dissociation constant (K_D)

Learn more about K_D

[Click here to learn more about \$K_D\$](#)

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab124956**).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-JNK1 + JNK2 + JNK3 (phospho T183+T183+T221) antibody [EPR5693] - BSA and Azide free (ab219584)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors