

Anti-Ki67 antibody [EPR3610] ab92742

敲除验证 重组 RabMAb


★★★★★ 12 Abreviews 312 References 17 图像

概述

产品名称	Anti-Ki67抗体[EPR3610]
描述	兔单克隆抗体[EPR3610] to Ki67
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), WB, IHC-P, ICC/IF
种属反应性	与反应: Human 不与反应: Mouse, Rat
免疫原	Synthetic peptide within Human Ki67 aa 1050-1150. The exact sequence is proprietary. Database link: P46013-1
阳性对照	WB: HeLa and ramos cell lysates. IHC-P: Human tonsil, colon, ovarian carcinoma, squamous cell carcinoma of cervix and colonic adenocarcinoma tissues. ICC/IF: HeLa, HT-29 cells, HAP1 cells. Flow Cyt (intra): Ramos cells, HAP1 cells.
常规说明	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
解离常数 (K _D)	K _D = 1.24 x 10 ⁻¹¹ M



[Learn more about K_D](#)

存储溶液	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol, 0.05% BSA
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR3610
同种型	IgG

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab92742于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		1/100 - 1/150. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/5000. Predicted molecular weight: 359 kDa. For unpurified use at 1/500 - 1/1000.
IHC-P	★★★★★ (5)	1/500 - 1/1000. See IHC antigen retrieval protocols . For unpurified use at 1/500 - 1/1000.
ICC/IF		Use a concentration of 0.5 - 1 µg/ml. If fixing cells in 4% PFA, it is recommended to permeabilized cells with 0.1% Triton-X for 5 min.

靶标

功能


Required to maintain individual mitotic chromosomes dispersed in the cytoplasm following nuclear envelope disassembly (PubMed:27362226). Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the chromosome surface (PubMed:27362226). Prevents chromosomes from collapsing into a single chromatin mass by forming a steric and electrostatic charge barrier: the protein has a high net electrical charge and acts as a surfactant, dispersing chromosomes and enabling independent chromosome motility (PubMed:27362226). Binds DNA, with a preference for supercoiled DNA and AT-rich DNA (PubMed:10878551). Does not contribute to the internal structure of mitotic chromosomes (By similarity). May play a role in chromatin organization (PubMed:24867636). It is however unclear whether it plays a direct role in chromatin organization or whether it is an indirect consequence of its function in maintaining mitotic chromosomes dispersed.

序列相似性

Contains 1 FHA domain.
Contains 16 K167R repeats.
Contains 1 PP1-binding domain.

发展阶段	Expression occurs preferentially during late G1, S, G2 and M phases of the cell cycle, while in cells in G0 phase the antigen cannot be detected (at protein level) (PubMed:6206131). Present at highest level in G2 phase and during mitosis (at protein level). In interphase, forms fiber-like structures in fibrillarin-deficient regions surrounding nucleoli (PubMed:2674163, PubMed:8799815).
翻译后修饰	Phosphorylated. Hyperphosphorylated in mitosis (PubMed:10502411, PubMed:10653604). Hyperphosphorylated form does not bind DNA.
细胞定位	Chromosome. Nucleus. Nucleus, nucleolus. Associates with the surface of the mitotic chromosome, the perichromosomal layer, and covers a substantial fraction of the mitotic chromosome surface (PubMed:27362226). Associates with satellite DNA in G1 phase (PubMed:9510506). Binds tightly to chromatin in interphase, chromatin-binding decreases in mitosis when it associates with the surface of the condensed chromosomes (PubMed:15896774, PubMed:22002106). Predominantly localized in the G1 phase in the perinucleolar region, in the later phases it is also detected throughout the nuclear interior, being predominantly localized in the nuclear matrix (PubMed:22002106).

图片




Flow cytometry overlay histogram showing wild-type Hap1 (green line) and MKI67 knockout Hap1 stained with ab92742 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab92742) (1×10^6 in 100µl at 0.04 µg/ml (1/57000)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C

Isotype control antibody Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control was used at the same concentration and conditions as the primary antibody (wild-type Hap1 - black line, MKI67 knockout Hap1 - grey line). Unlabelled sample was also used as a control (this line is not shown for the purpose of simplicity).

Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.

This antibody gave a positive signal in Hap1 Fixed with 4% formaldehyde (10 min) / permeabilised with 0.1% PBS-Triton X-100 for 15 min under the same conditions.




Comparison between RNAPII-S2P-low cells and Ki-67- cells

a: Regulation of Ki-67 and RNAPII-S2P during proliferation and quiescence in T98G glioblastoma cells. T98G cells were grown in culture medium containing 10% (v/v) fetal bovine serum (FBS), were induced to become quiescent by serum starvation in medium supplemented with 0.5% (v/v) FBS for 14 days, and then were re-stimulated by being split 1:5 into new medium containing 10% (v/v) FBS and cultured for 3 days. The cells were detached from dishes with trypsin-EDTA solution, fixed in 10% (v/v) neutral buffered formalin, and centrifuged. Paraffin sections of the pellet were cut, and expression of Ki-67 and RNAPII-S2P was examined by single (brown; a1, a2, a4, a5, a7, a8) or double immunostaining (Ki-67, brown; RNAPII-S2P, red; a3, a6, a9). Hematoxylin (blue) was used as a nuclear stain. Ki-67- RNAPII-S2P-low cells (blue cells in the double stained sections) emerged only in the quiescent condition (a6, arrows). Scale bar, 10 μ m. b: Single-color immunostaining for Ki-67 (b1) and RNAPII-S2P (b2) in serial sections of glioblastoma tissue. Ki-67- tumor cells were frequently found, whereas only a few RNAPII-S2P-low cells (arrows) were observed around necrotic area. N, necrotic area; V, blood vessels. Scale bars, 50 μ m.


Ki67 detected using ab92742.

(From Figure S2 of Ishii et al)



ab92742 staining Ki67 in wild-type HAP1 cells (top panel) and Ki67 knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab92742 at 1 μ g/ml and **ab195889** at 1/250 dilution (shown in pseudo colour red) overnight at +4°C, followed by a further incubation at room temperature for 1h with a goat secondary antibody to Rabbit IgG (Alexa Fluor® 488) (**ab150081**) at 2 μ g/ml (shown in green). Nuclear DNA was labeled in blue with DAPI.

Alexa Fluor® 488 (**ab197234**) and Alexa Fluor® 647 (**ab196907**) conjugated versions are available for this clone.



Anti-Ki67 antibody [EPR3610] (ab92742) at 1/5000 dilution (purified) + Ramos (Human Burkitt's lymphoma cell line) cell lysate at 20 µg


Secondary

Peroxidase-conjugated goat anti-rabbit IgG, (H+L) at 1/1000 dilution

Predicted band size: 359 kDa

Observed band size: 395 kDa


Blocking and dilution buffer: 5% NFDM/TBST.



Overlay histogram showing Ramos (Human Burkitt's lymphoma cell line) cells stained with unpurified ab92742 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (unpurified ab92742, 1/100 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) ([ab96899](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (1 µg/1x10⁶ cells) used under the same conditions.

Acquisition of >5,000 events was performed.


Alexa Fluor®488 ([ab197234](#)) and Alexa Fluor®647 ([ab196907](#)) conjugated versions are available for this clone.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human squamous cell carcinoma of cervix tissue labeling Ki67 with purified ab92742 at 1/250. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9. **ab97051**, a HRP-conjugated goat anti-rabbit IgG (H+L) was used as the secondary antibody (1/500). Counterstained with hematoxylin.

Negative control using PBS instead of primary antibody (inset).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610] (ab92742)

Chromogenic triple immunostaining for estrogen receptor (ER), progesterone receptor (PgR), and Ki-67 in breast cancer tissue to verify the triple immunostaining detection method.

Panel e: ER⁺ PgR⁻ Ki-67⁻ cells were stained red (short arrow), ER⁻ PgR⁺ Ki-67⁻ cells were stained blue (black arrowhead), ER⁺ PgR⁺ Ki-67⁻ cells were stained purple (long arrow), and Ki-67⁺ cells were stained brown (white arrowhead). These colors are easily distinguishable. Scale bars, 25 μ m.

Deparaffinized sections were pretreated for antigen retrieval by boiling in antigen retrieval solution, pH 9. Sections were incubated with rabbit monoclonal antibody against Ki67 ab92742 at a 1/1000 dilution. After the reaction with (HRP)-conjugated secondary antibodies color was developed with (DAB) and sections were counterstained with hematoxylin.

Ishii et al PLoS One. 2016 Jan 22;11(1):e0147366. doi: 10.1371/journal.pone.0147366. eCollection 2016. Fig S4. Reproduced under the Creative Commons license <http://creativecommons.org/licenses/by/4.0/>




Immunocytochemistry/ Immunofluorescence - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunocytochemistry analysis of HT-29 (Human colorectal adenocarcinoma cell line) cells labeling Ki67 with purified ab92742 at 1/250. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500) was used as the secondary antibody. DAPI (blue) was used as the nuclear counterstain. **ab7291**, a mouse anti-tubulin (1/1000) and **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500) were also used.


Control 1: primary antibody (1/250) and secondary antibody, **ab150120**, an Alexa Fluor® 594-conjugated goat anti-mouse IgG (1/500).

Control 2: **ab7291** (1/1000) and secondary antibody, **ab150077**, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/500).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human normal colon tissue labeling Ki67 with unpurified ab92742. Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human colonic adenocarcinoma tissue labeling Ki67 with unpurified ab92742. Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.




Flow Cytometry (Intracellular) - Anti-Ki67 antibody
[EPR3610] (ab92742)

Intracellular Flow Cytometry analysis of Ramos (Human Burkitt's lymphoma cell line) cells lablling Ki67 with purified ab92742 at 1/150 (red). Cells were fixed with 2% paraformaldehyde. An FITC-conjugated goat anti-rabbit IgG (1/150) was used as the secondary antibody. Black - Isotype control, rabbit monoclonal IgG. Blue - Unlabeled control, cells without incubation with primary and secondary antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610]
(ab92742)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human ovarian carcinoma tissue labeling Ki67 with unpurified ab92742. Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Western blot - Anti-Ki67 antibody [EPR3610]
(ab92742)


Anti-Ki67 antibody [EPR3610] (ab92742) at 1/500 dilution (unpurified) + HeLa (Human epithelial cell line from cervix adenocarcinoma) cell lysate at 10 µg

Secondary

HRP-conjugated goat anti-rabbit IgG at 1/2000 dilution


Predicted band size: 359 kDa

Observed band size: 395 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human cervical carcinoma tissue labeling Ki67 with unpurified ab92742. Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-Ki67 antibody [EPR3610] (ab92742)

This image is courtesy of an anonymous Abreview.

ab92742 staining Ki67 in human adenocarcinoma cells by ICC (Immunocytochemistry).

Cells were fixed with paraformaldehyde and permeabilized with 0.1% Triton X-100 in PBS and blocked with 5% serum for 1 hour at 21°C. Samples were incubated with primary antibody (1/1000) for 12 hours at 4°C. A Cy3[®] conjugated donkey anti-rabbit IgG polyclonal was used as the secondary antibody at a dilution of 1/200.



Immunocytochemistry/ Immunofluorescence - Anti-Ki67 antibody [EPR3610] (ab92742)

Immunocytochemistry analysis of HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Ki67 with unpurified ab92742 at a dilution of 1/250.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Ki67 antibody [EPR3610] (ab92742)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors