

Anti-L Glutamate antibody ab62668

★★★★★ [1 Abreviews](#) [1 References](#)

概述

产品名称	Anti-L Glutamate抗体
描述	鸡多克隆抗体 to L Glutamate
宿主	Chicken
特异性	No measurable cross-reactivity (<1:1000) against glutamate in peptides or proteins. Fixed tissue cross-reactivity tested with known targets at recommended dilution. No measurable glutaraldehyde-fixed tissue cross-reactivity (<1:1000) against L-alanine, gamma-aminobutyrate, D/L-aspartate, 1-amino-4-guanidobutane (AGB), D/L-arginine, L-citrulline, L-cysteine, D/L-glutamine, glutathione, glycine, L-lysine, L-ornithine, L-serine, taurine, L-threonine, L-tryptophan, L-tyrosine. Modest cross-reactivity (1:20) against D-glutamate. Significant cross-reactivity (1:8) against free NAAG in competition assays (NAAG is not retained in aldehyde-fixed tissue).
经测试应用	适用于: ICC
种属反应性	与反应: Species independent
免疫原	L Glutamate cross-linked to purified fraction V bovine serum albumin with glutaraldehyde.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.05% Thimerosal (merthiolate) Constituents: 0.268% PBS, 1% Normal Goat Serum
纯度	IgY fraction
纯化说明	IgYs were separated from other yolk proteins by serial PEG6000 extractions using a modification of the method of Polsonet al Immunological Investigations. 1985, 14, 323-327.

克隆 多克隆
同种型 IgY

应用

The Abpromise guarantee **Abpromise™**承诺保证使用ab62668于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC		1/100. Endogenous content mapping by LM and EM immunocytochemistry. True dilution at user dilution: 1/32000. Optimal fixation: 0.1-2.5% glutaraldehyde, 1% formaldehyde using HPI (High Performance Immunocytochemistry). Minimum glutaraldehyde: 0.0% using EHPI (Enhanced HPI) with 4% formaldehyde.

靶标

相关性 L Glutamate is a non-essential amino acid naturally occurring in the L-form. It is the most common excitatory neurotransmitter in the Central Nervous System.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors