

Anti-LDL Receptor antibody ab30532

★★★★★ [12 Abreviews](#) [82 References](#) [6 图像](#)

概述

产品名称	Anti-LDL Receptor抗体
描述	兔多克隆抗体to LDL Receptor
宿主	Rabbit
经测试应用	适用于: IHC-P, ICC/IF, WB
种属反应性	与反应: Mouse, Rat, Human
免疫原	Synthetic peptide: SVADTKGVKRRTL , corresponding to amino acids 499-511 of Mouse LDL Receptor Run BLAST with Run BLAST with

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
存储溶液	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: Tris buffered saline, 40% Glycerol, 0.1% BSA
纯度	Immunogen affinity purified
克隆	多克隆
同种型	IgG

应用

The Abpromise guarantee


Abpromise™承诺保证使用ab30532于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-P	★★★★★ (1)	1/400.
ICC/IF	★★★★★ (1)	1/100 - 1/400.
WB	★★★★☆ (5)	Use at an assay dependent concentration. Predicted molecular weight: 95 kDa.

靶标	
功能	Binds LDL, the major cholesterol-carrying lipoprotein of plasma, and transports it into cells by endocytosis. In order to be internalized, the receptor-ligand complexes must first cluster into clathrin-coated pits. In case of HIV-1 infection, functions as a receptor for extracellular Tat in neurons, mediating its internalization in uninfected cells.
疾病相关	Defects in LDLR are the cause of familial hypercholesterolemia (FH) [MIM:143890]; a common autosomal semi-dominant disease that affects about 1 in 500 individuals. The receptor defect impairs the catabolism of LDL, and the resultant elevation in plasma LDL-cholesterol promotes deposition of cholesterol in the skin (xanthelasma), tendons (xanthomas), and coronary arteries (atherosclerosis).
序列相似性	Belongs to the LDLR family. Contains 3 EGF-like domains. Contains 7 LDL-receptor class A domains. Contains 6 LDL-receptor class B repeats.
翻译后修饰	N- and O-glycosylated. Ubiquitinated by MYLIP leading to degradation.
细胞定位	Cell membrane. Endomembrane system. Membrane > clathrin-coated pit. Found distributed from the plasma membrane to intracellular compartments.


图片



Immunocytochemistry - Anti-LDL Receptor antibody
(ab30532)

This image is courtesy of an anonymous Abreview

Immunocytochemistry analysis of acetone-fixed human Liver cells, Hep2G staining with ab30532 at 1/200 dilution. Secondary antibody was FITC rabbit IgG at 1/400 dilution. Samples were incubated with the primary antibody with Serum/PBS for 1 hours at 21°C. Blocking was done using 5% serum for 15 minutes at 21°C.



Western blot - Anti-LDL Receptor antibody
(ab30532)

All lanes : Anti-LDL Receptor antibody (ab30532) at 1/200 dilution

Lane 1 : RAW 264.7 Cells at 25 µg

Lane 2 : RAW 264.7 Cells at 50 µg


Lane 3 : Rat Liver Supernatant at 50 µg

Secondary

All lanes : Anti-Rabbit-IgG-HRP at 1/7000 dilution


Developed using the ECL technique.

Predicted band size: 95 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LDL Receptor antibody (ab30532)

Immunohistochemistry analysis of formalin-fixed, paraffin-embedded (FFPE) mouse lung tissue after heat induced antigen retrieval in pH 6.0 citrate buffer. After incubation with LDL receptor polyclonal antibody, ab30532, at a 1:100 dilution, slides were incubated with biotinylated secondary antibody, followed by alkaline phosphatase-streptavidin and chromogen (DAB).



Western blot - Anti-LDL Receptor antibody (ab30532)

All lanes : Anti-LDL Receptor antibody (ab30532)

Lane 1 : HeLa Cell Lysate

Lane 2 : HepG2 Cell Lysate


Lane 3 : Huh7 Cell Lysate

Lysates/proteins at 50 µg per lane.

Secondary

All lanes : Mouse monoclonal [6F12] to Cytomegalovirus ([ab6500](#)) (Anti-Rabbit IgG (HRP) conjugate)

Predicted band size: 95 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-LDL Receptor antibody (ab30532)

ab30532 staining LDL receptor in human lung parenchyma.

Left panel: with primary antibody at 1/400. Right panel: isotype control.

Sections were stained using an automated system (DAKO Autostainer Plus), at room temperature: sections were rehydrated and antigen retrieved with the Dako 3 in 1 AR buffers citrate pH6.1. Slides were peroxidase blocked in 3% H₂O₂ in methanol for 10 mins. They were then blocked with Dako Protein block for 10 minutes (containing casein 0.25% in PBS) then incubated with primary antibody for 20 min and detected with Dako envision flex amplification kit for 30 minutes. Colorimetric detection was completed with Diaminobenzidine for 5 minutes. Slides were counterstained with Haematoxylin and coverslipped under DePeX.

Please note that for manual staining we recommend to optimize the primary antibody concentration and incubation time (overnight incubation), and amplification may be required.



Immunocytochemistry/ Immunofluorescence - Anti-LDL Receptor antibody (ab30532)

ICC/IF image of ab30532 stained HepG2 cells. The cells were 4% PFA fixed (10 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab30532, 1/400) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors