

# Anti-Macrophage Inflammatory Protein 3 alpha antibody ab106151

[2 References](#) [1 图像](#)

### 概述

<b>产品名称</b>	Anti-Macrophage Inflammatory蛋白3 alpha抗体
<b>描述</b>	兔多克隆抗体to Macrophage Inflammatory蛋白3 alpha
<b>宿主</b>	Rabbit
<b>经测试应用</b>	<b>适用于:</b> WB
<b>种属反应性</b>	<b>与反应:</b> Mouse
<b>免疫原</b>	ab106151 was prepared from whole Rabbit serum produced by repeated immunizations with recombinant full length Mouse MIP-3a protein.
<b>常规说明</b>	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 性能

<b>形式</b>	Lyophilized:Centrifuge product if not completely clear after standing at room temperature. Restore with 0.1 mL of deionized water (or equivalent). Dilute only prior to immediate use. Expiration date is one (1) year from date of restoration.
<b>存放说明</b>	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
<b>存储溶液</b>	Constituents: 0.88% Sodium chloride, 0.42% Potassium phosphate
<b>纯度</b>	Protein A purified
<b>克隆</b>	多克隆
<b>同种型</b>	IgG

### 应用

## The Abpromise guarantee

**Abpromise™**承诺保证使用ab106151于以下的经测试应用


“应用说明”部分下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/1000. Predicted molecular weight: 11 kDa.

## 靶标

<b>功能</b>	Chemotactic factor that attracts lymphocytes and, slightly, neutrophils, but not monocytes. Inhibits proliferation of myeloid progenitors in colony formation assays. May be involved in formation and function of the mucosal lymphoid tissues by attracting lymphocytes and dendritic cells towards epithelial cells. C-terminal processed forms have been shown to be equally chemotactically active for leukocytes. Possesses antibacterial activity E.coli ATCC 25922 and S.aureus ATCC 29213.
<b>组织特异性</b>	Expressed predominantly in the liver, lymph nodes, appendix, peripheral blood lymphocytes, and fetal lung. Low levels seen in thymus, prostate, testis, small intestine and colon.
<b>序列相似性</b>	Belongs to the intercrine beta (chemokine CC) family.
<b>翻译后修饰</b>	C-terminal processed forms which lack 1, 3 or 6 amino acids are produced by proteolytic cleavage after secretion from peripheral blood monocytes.
<b>细胞定位</b>	Secreted.

## 图片



Anti-Macrophage Inflammatory Protein 3 alpha antibody (ab106151) at 1/1000 dilution + Mouse recombinant MIP3a protein (the immunogen) at 0.25 µg

### Secondary

HRP Goat anti-Rabbit IgG at 1/40000 dilution

**Predicted band size:** 11 kDa

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery

- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors