abcam

Product datasheet

Anti-MCP5 antibody ab9737

概述

产品名称 Anti-MCP5抗体

描述 兔多克隆抗体to MCP5

宿主 Rabbit

经测试应用 适用于: ELISA, Sandwich ELISA, WB, Neutralising

种属反应性 与反应: Mouse

免疫原 Recombinant full length protein corresponding to Mouse MCP5. Highly pure (>98%) recombinant

mMCP-5 (mouse Macrophage/Monocyte Chemotactic Protein-5

Database link: Q62401

阳性对照 Recombinant mouse MCP5 protein (<u>ab9738</u>) can be used as a positive control in WB.

常规说明

The Life Science industry has been in the grips of a reproducibility crisis for a number of years.

Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets

your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be

found below, along with publications, customer reviews and Q&As

性能

形式 Lyophilized:Reconstitute with 200µl of sterile water.

存放说明 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term.

unknown

存储溶液 No preservative, sterile filtered

纯**度** Immunogen affinity purified

克隆多克隆同种型unknown

应用

轻链类型

The Abpromise guarantee Abpromise Abpromise Guarantee

Abpromise™承诺保证使用ab9737于以下的经测试应用

1

"应用说明"部分 下显示的仅为推荐的起始稀释度:实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ELISA		Use at an assay dependent concentration.
Sandwich ELISA		Use at an assay dependent concentration. To detect mMCP-5 by sandwich ELISA (using 100 ul/well antibody solution) a concentration of 0.5 - 2.0 ug/ml of this antibody is required.
WB		Use at an assay dependent concentration.
Neutralising		Use at an assay dependent concentration.

12.00		
功能	Chemotactic factor that attracts eosinophils, monocytes, and lymphocytes but not neutrophils. Potent monocyte active chemokine that signals through CCR2. Involved in allergic inflammation and the host response to pathogens and may play a pivotal role during early stages of allergic lung inflammation.	
组织 特异性	Predominantly expressed in the lymph nodes and thymus. Also found in the salivary glands containing lymph nodes, breast, heart, lung, brain, small intestine, kidney and colon.	
序列相似性	Belongs to the intercrine beta (chemokine CC) family.	
细胞定位	Secreted.	

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.cn/abpromise or contact our technical team.

Terms and conditions

如卡

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors