


Anti-MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211) antibody ab4759

2 图像

概述

产品名称	Anti-MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211)抗体
描述	兔多克隆抗体to MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211)
宿主	Rabbit
特异性	This antibody reacts with MEK 3 and MEK 6 (93% homologous). This antibody shows some cross-reactivity with MEK 4 when tested in a system with high MEK 4 expression levels. It does not react with any other MEK isoforms.
经测试应用	适用于: WB
种属反应性	与反应: Human 预测可用于: a wide range of other species 
免疫原	Synthetic peptide corresponding to Human MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211).
阳性对照	Mal-E-tagged fusion protein expressing MEK 6, left inactivated or activated by adding MEKK, Anisomycin treated A549 cell line
常规说明	<p>Mitogen Activated Protein Kinase Kinases 3 and 6 (MEK 3/6 or MKK 3/6) are 42 kDa members of a tyrosine/threonine protein kinase family that activate p38, which is part of the inflammation/stress signaling pathway. Phosphorylation of MEK3 and 6 by MEKK1 on serine 189 and threonine 193 (serine 207 and threonine 211 for MEK6) in the catalytic domain activates the proteins and enables them to phosphorylate p38.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw

	cycles.
存储溶液	<p>pH: 7.30</p> <p>Preservative: 0.05% Sodium azide</p> <p>Constituents: PBS, 0.1% BSA</p>
	BSA is IgG and protease free
纯度	Immunogen affinity purified
纯化说明	Purified from rabbit serum by sequential epitope-specific chromatography. The antibody has been negatively preadsorbed using a non-phosphopeptide corresponding to the site of phosphorylation to remove antibody that is reactive with non-phosphorylated MEK 3. The final product is generated by affinity chromatography using a MEK 3 derived peptide that is phosphorylated at serine 189 and threonine 193.
Primary antibody说明	Mitogen Activated Protein Kinase Kinases 3 and 6 (MEK 3/6 or MKK 3/6) are 42 kDa members of a tyrosine/threonine protein kinase family that activate p38, which is part of the inflammation/stress signaling pathway. Phosphorylation of MEK3 and 6 by MEKK1 on serine 189 and threonine 193 (serine 207 and threonine 211 for MEK6) in the catalytic domain activates the proteins and enables them to phosphorylate p38.
克隆	多克隆
同种型	IgG

应用


The Abpromise guarantee Abpromise™承诺保证使用ab4759于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
WB		1/500 - 1/2000. Predicted molecular weight: 39.3 kDa.

靶标

图片



All lanes : Anti-MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211) antibody (ab4759) at 1/2000 dilution

Lane 1 : A549 whole cell extract

Lane 2 : A549 treated for 20 minutes with 200 nM of PMA

Lane 3 : A549 treated for 30 minutes with 25 µg/mL of Anisomycin

Lane 4 : A549 treated for 40 minutes with UV

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat anti-Rabbit IgG (H+L) Secondary Antibody, HRP conjugate at 1/2500 dilution

Predicted band size: 39.3 kDa

Observed band size: 39, 37 kDa

Two bands ~ 37 and 39 kDa corresponding to MEK3 (pS189/pT193) + MEK6 (pS207/pT211) was observed in Anisomycin treated cell line tested



Western blot - Anti-MEK3 (phospho S189/T193) + MEK6 (phospho S207/T211) antibody (ab4759)

Peptide Competition: Extracts prepared from background extracts with Mal-E tagged fusion protein expressing MEK 6 left inactivated (6) or with MEKK added for activation (1-5) were resolved by SDS-PAGE on a 10% polyacrylamide gel and transferred to PVDF. Membranes were blocked with a 5% BSA-TBST buffer overnight at 4°C, then were incubated with 0.50 µg/mL ab4759 antibody for two hours at room temperature in a 3% BSA-TBST buffer, following prior incubation with: no peptide (1, 6), the non-phosphopeptide corresponding to the phosphopeptide immunogen (2), a generic phosphoserine-containing peptide (3), a generic phosphothreonine-containing peptide (4), or the phosphopeptide immunogen (5). After washing, the membrane was incubated with goat F(ab')₂ anti-rabbit IgG alkaline phosphatase and signals were detected. The data show that only the phosphopeptide corresponding to MEK3/6 [pSpT189/193]/[pSpT207/211] blocks the antibody signal, demonstrating the specificity of the antibody. Please note that the Mal-E-tagged fusion protein expressing MEK6 runs at ~81 kDa.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors