

Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free ab222925

敲除验证
重组
RabMAb

10 图像

概述

产品名称	Anti-Met (c-Met)抗体[EPR19067] - Low endotoxin, Azide free
描述	兔单克隆抗体[EPR19067] to Met (c-Met) - Low endotoxin, Azide free
宿主	Rabbit
经测试应用	适用于: Flow Cyt (Intra), IHC-P, ICC/IF, WB, Indirect ELISA
种属反应性	与反应: Human, Recombinant fragment
免疫原	Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: A549, HeLa and HepG2 whole cell lysates; Human liver lysate; 293T whole cell lysate transfected with a His-tagged human c-Met construct; HeLa whole cell lysate, untreated or treated with PNGase F. IHC-P: Human breast, colon, liver cancer and ovary cancer tissues. ICC/IF: HeLa and A549 cells. Flow Cyt (intra): A549 and HeLa cells.
常规说明	ab222925 is the carrier-free version of ab216574 .

Our **carrier-free** antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.

This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our **conjugation kits** for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit

monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Our **Low endotoxin, azide-free formats** have low endotoxin level (≤ 1 EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.2 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR19067
同种型	IgG

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab222925于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent concentration.
WB		Use at an assay dependent concentration. Detects a band of approximately 45-175 kDa (predicted molecular weight: 155 kDa).
Indirect ELISA		Use at an assay dependent concentration.

靶标

功能	Receptor for hepatocyte growth factor and scatter factor. Has a tyrosine-protein kinase activity. Functions in cell proliferation, scattering, morphogenesis and survival.
疾病相关	Note=Activation of MET after rearrangement with the TPR gene produces an oncogenic protein.

Note=Defects in MET may be associated with gastric cancer.

Defects in MET are a cause of hepatocellular carcinoma (HCC) [MIM:114550].

Defects in MET are a cause of renal cell carcinoma papillary (RCCP) [MIM:605074]. It is a subtype of renal cell carcinoma tending to show a tubulo-papillary architecture formed by numerous, irregular, finger-like projections of connective tissue. Renal cell carcinoma is a heterogeneous group of sporadic or hereditary carcinoma derived from cells of the proximal renal tubular epithelium. It is subclassified into common renal cell carcinoma (clear cell, non-papillary carcinoma), papillary renal cell carcinoma, chromophobe renal cell carcinoma, collecting duct carcinoma with medullary carcinoma of the kidney, and unclassified renal cell carcinoma.

Note=A common allele in the promoter region of the MET shows genetic association with susceptibility to autism in some families. Functional assays indicate a decrease in MET promoter activity and altered binding of specific transcription factor complexes.

Note=MET activating mutations may be involved in the development of a highly malignant, metastatic syndrome known as cancer of unknown primary origin (CUP) or primary occult malignancy. Systemic neoplastic spread is generally a late event in cancer progression. However, in some instances, distant dissemination arises at a very early stage, so that metastases reach clinical relevance before primary lesions. Sometimes, the primary lesions cannot be identified in spite of the progresses in the diagnosis of malignancies.

序列相似性

Belongs to the protein kinase superfamily. Tyr protein kinase family.

Contains 3 IPT/TIG domains.

Contains 1 protein kinase domain.

Contains 1 Sema domain.

结构域

The kinase domain is involved in SPSB1 binding.


翻译后修饰

Dephosphorylated by PTPRJ at Tyr-1349 and Tyr-1365.

细胞定位

Membrane.

图片



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunohistochemical analysis of paraffin-embedded human breast tissue labeling Met (c-Met) with [ab216574](#) at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Membranous staining on human breast is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab216574](#)).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunohistochemical analysis of paraffin-embedded human colon tissue labeling Met (c-Met) with **ab216574** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Membranous staining on human colon is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunohistochemical analysis of paraffin-embedded human liver cancer tissue labeling Met (c-Met) with **ab216574** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Membranous staining on tumor cells of human liver cancer is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue labeling Met (c-Met) with **ab216574** at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Cytoplasmic and membranous staining on tumor cells of human ovary cancer is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Met (c-Met) with **ab216574** at 1/1000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) secondary antibody at 1/1000 dilution (green).


Confocal image showing cytoplasmic staining on HeLa cell line.

The nuclear counterstain is DAPI (blue).

Tubulin is detected with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor[®] 488) (**ab150077**) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).



Immunocytochemistry/ Immunofluorescence - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized A549 (Human lung carcinoma cell line) cells labeling Met (c-Met) with **ab216574** at 1/1000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) secondary antibody at 1/1000 dilution (green).


Confocal image showing cytoplasmic staining on A549 cell line.

The nuclear counterstain is DAPI (blue).

Tubulin is detected with **ab195889** (Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594)) at 1/200 dilution (red).

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat anti-rabbit IgG (Alexa Fluor® 488) (**ab150077**) at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).




Flow Cytometry (Intracellular) - Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed A549 (Human lung carcinoma cell line) cells labeling Met (c-Met) with **ab216574** at 1/600 dilution (red) compared with a rabbit monoclonal IgG isotype control (**ab172730**; black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab216574**).


Indirect ELISA antibody dose-response curve antigen at 1000 ng/ml



Indirect ELISA - Anti-Met (c-Met) antibody
[EPR19067] - Low endotoxin, Azide free (ab222925)

This data was developed using [ab216574](#), the same antibody clone in a different buffer formulation.

ELISA analysis of Human c-met recombinant protein at 1000 ng/mL with [ab216574](#). An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) at 1/2500 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-Met (c-Met)
antibody [EPR19067] - Low endotoxin, Azide free
(ab222925)

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling Met (c-Met) with [ab216574](#) at 1/600 dilution (red) compared with a rabbit monoclonal IgG isotype control ([ab172730](#); black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab216574](#)).

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Met (c-Met) antibody [EPR19067] - Low endotoxin, Azide free (ab222925)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors