

# Anti-MMP2 antibody [EPR1184] - Low endotoxin, Azide free ab215986

**重组** RabMAb

**24 References** [6 图像](#)

### 概述

产品名称	Anti-MMP2抗体[EPR1184] - Low endotoxin, Azide free
描述	兔单克隆抗体[EPR1184] to MMP2 - Low endotoxin, Azide free
宿主	Rabbit
特异性	<p>In Western Blot, this product typically gives a weaker signal in some hepatocarcinoma like HepG2, MHCC97L(PMID:33184263 ) et al. Please use recommended positive control when testing these cells.</p> <p>Compared with <a href="#">ab92536</a>, <a href="#">ab181286</a> has higher sensitivity. We recommend <a href="#">ab181286</a> as an alternative for testing MMP2 in western blot.</p>
经测试应用	<p><b>适用于:</b> Flow Cyt (Intra), WB, ICC/IF</p> <p><b>不适用于:</b> IHC-P</p>
种属反应性	<b>与反应:</b> Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
阳性对照	WB: HepG2, L6, Raw264.7 and NIH/3T3 cell lysates; fetal heart and human skin tissue lysate; Human plasma, brain and breast tissue lysate ICC/IF: PC-3 cells Flow Cyt (intra): HeLa and PC-3 cells
常规说明	<p>ab215986 is the carrier-free version of <a href="#">ab92536</a>.</p> <p>Our <b>carrier-free</b> antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our <a href="#">conjugation kits</a> for antibody conjugates that are ready-to-use in as little as 20 minutes with &lt;1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar<sup>®</sup> Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar<sup>®</sup> is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p>

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information [see here](#).

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to [RabMAb® patents](#).

Our **Low endotoxin, azide-free formats** have low endotoxin level ( $\leq 1$  EU/ml, determined by the LAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

## 性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C. Do Not Freeze.
存储溶液	pH: 7.20 Constituent: PBS
无载体	是
纯度	Protein A purified
克隆	单克隆
克隆编号	EPR1184
同种型	IgG

## 应用

**The Abpromise guarantee** [Abpromise™](#) 承诺保证使用 ab215986 于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度; 实际最佳的稀释度/浓度应由使用者检定。

应用	Ab 评论	说明
Flow Cyt (Intra)		Use at an assay dependent concentration. <b>ab199376</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Compared with <b>ab92536</b> , <b>ab181286</b> has higher sensitivity. We recommend <b>ab181286</b> as an alternative for testing MMP2 in western blot.
ICC/IF		Use at an assay dependent concentration.

应用说明 Is unsuitable for IHC-P.

## 靶标

功能 Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the

vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-

-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro.

PEX, the C-terminal non-catalytic fragment of MMP2, possesses anti-angiogenic and anti-tumor properties and inhibits cell migration and cell adhesion to FGF2 and vitronectin. Ligand for integrin/beta3 on the surface of blood vessels.

#### 组织特异性

Produced by normal skin fibroblasts. PEX is expressed in a number of tumors including gliomas, breast and prostate.

#### 疾病相关

Defects in MMP2 are the cause of Torg-Winchester syndrome (TWS) [MIM:259600]; also known as multicentric osteolysis nodulosis and arthropathy (MONA). TWS is an autosomal recessive osteolysis syndrome. It is severe with generalized osteolysis and osteopenia. Subcutaneous nodules are usually absent. Torg-Winchester syndrome has been associated with a number of additional features including coarse face, corneal opacities, patches of thickened, hyperpigmented skin, hypertrichosis and gum hypertrophy. However, these features are not always present and have occasionally been observed in other osteolysis syndromes.

#### 序列相似性

Belongs to the peptidase M10A family.

Contains 3 fibronectin type-II domains.

Contains 4 hemopexin-like domains.

#### 结构域

The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.


#### 翻译后修饰

Phosphorylation on multiple sites modulates enzymatic activity. Phosphorylated by PKC in vitro. The propeptide is processed by MMP14 (MT-MMP1) and MMP16 (MT-MMP3). Autocatalytic cleavage in the C-terminal produces the anti-angiogenic peptide, PEX. This processing appears to be facilitated by binding integrin/beta3.

#### 细胞定位

Secreted > extracellular space > extracellular matrix. Membrane. Nucleus. Colocalizes with integrin alphaV/beta3 at the membrane surface in angiogenic blood vessels and melanomas. Found in mitochondria, along microfibrils, and in nuclei of cardiomyocytes.

#### 图片



Western blot - Anti-MMP2 antibody [EPR1184] -  
Low endotoxin, Azide free (ab215986)

**All lanes :** Anti-MMP2 antibody [EPR1184] (**ab92536**) at 1/1000 dilution

**Lane 1 :** Human plasma tissue lysate

**Lane 2 :** Human brain tissue lysate

**Lane 3 :** Human breast tissue lysate

Lysates/proteins at 20 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

**Observed band size:** 69,72 kDa


**Exposure time:** 60 seconds

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab92536**).

Blocking and diluting buffer and concentration: 5% NFDM/TBST.

**ab181602** was used as a GAPDH loading control.

Compared with **ab92536**, **ab181286** has higher sensitivity. We recommend **ab181286** as an alternative for testing MMP2 in western blot.



Western blot - Anti-MMP2 antibody [EPR1184] -  
Low endotoxin, Azide free (ab215986)

**All lanes :** Anti-MMP2 antibody [EPR1184] (**ab92536**) at 1/1000 dilution

**Lane 1 :** HT-1080 (Human fibrosarcoma epithelial cell) whole cell lysate

**Lane 2 :** Untreated HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

**Lane 3 :** HepG2 (Human hepatocellular carcinoma epithelial cell) treated with 300ng/ml BFA for 24 hours whole cell lysate

**Lane 4 :** Huh7 (Human hepatocellular carcinoma epithelial cell) whole cell lysate

**Lane 5 :** U-87 MG (Human glioblastoma-astrocytoma epithelial cell) whole cell lysate

Lysates/proteins at 20 µg per lane.

## Secondary

**All lanes :** Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

**Observed band size:** 69,72 kDa


**Exposure time:** 60 seconds

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92536](#)).

Blocking and diluting buffer and concentration: 5% NFDM /TBST.

[ab181602](#) was used as GAPDH loading control.


Compared with [ab92536](#), [ab181286](#) has higher sensitivity. We recommend [ab181286](#) as an alternative for testing MMP2 in western blot.



Flow Cytometry (Intracellular) - Anti-MMP2 antibody  
[EPR1184] - Low endotoxin, Azide free (ab215986)


Intracellular Flow Cytometry analysis of PC-3 (human prostate adenocarcinoma) cells labeling MMP2 with purified [ab92536](#) at 1/180 dilution (10ug/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. A Goat anti rabbit IgG (Alexa Fluor® 488) (1/2000 dilution) was used as the secondary antibody. Rabbit monoclonal IgG (Black) was used as the isotype control, cells without incubation with primary antibody and secondary antibody (Blue) were used as the unlabeled control.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide ([ab92536](#)).



Flow Cytometry (Intracellular) - Anti-MMP2 antibody  
[EPR1184] - Low endotoxin, Azide free (ab215986)

Overlay histogram showing HeLa cells fixed in 4% PFA and stained with purified **ab92536** at a dilution of 1 in 400 (red line). The secondary antibody used was FITC goat anti-rabbit at a dilution of 1 in 500. Rabbit monoclonal IgG was used as an isotype control (black line) and cells incubated in the absence of both primary and secondary antibody were used as a negative control (blue line). This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab92536**).



Immunocytochemistry/ Immunofluorescence - Anti-MMP2 antibody [EPR1184] - Low endotoxin, Azide free (ab215986)

Immunofluorescence staining of PC-3 cells with purified **ab92536** at a working dilution of 1/250, counter-stained with DAPI. The secondary antibody was Alexa Fluor<sup>®</sup> 488 goat anti-rabbit (**ab150077**), used at a dilution of 1/1000. **ab7291**, a mouse anti-tubulin antibody (1/1000), was used to stain tubulin along with **ab150120** (Alexa Fluor<sup>®</sup> 594 goat anti-mouse, 1/1000), shown in the top right hand panel. The cells were fixed in 4% PFA and permeabilized using 0.1% Triton X 100. The negative controls are shown in bottom middle and right hand panels - for negative control 1, purified **ab92536** was used at a dilution of 1/500 followed by an Alexa Fluor<sup>®</sup> 594 goat anti-mouse antibody (**ab150120**) at a dilution of 1/500. For negative control 2, **ab7291** (mouse anti-tubulin) was used at a dilution of 1/500 followed by an Alexa Fluor<sup>®</sup> 488 goat anti-rabbit antibody (**ab150077**) at a dilution of 1/400.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (**ab92536**).

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-MMP2 antibody [EPR1184] - Low endotoxin,  
Azide free (ab215986)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors