

Anti-Monocyte + Macrophage antibody [MOMA-2] ab33451

★★★★★ **3 Abreviews** **92 References**

概述

产品名称	Anti-Monocyte + Macrophage抗体[MOMA-2]
描述	大鼠单克隆抗体[MOMA-2] to Monocyte + Macrophage
宿主	Rat
特异性	This antibody recognises an intracellular antigen of mouse macrophages and monocytes. It reacts strongly with macrophages in lymphoid organs in all mouse strains tested. Staining may also be seen on the epithelium of mucosa.
经测试应用	适用于: IHC-Fr, IHC-P, IHC-FoFr, Flow Cyt
种属反应性	与反应: Mouse
免疫原	Tissue, cells or virus corresponding to Mouse Monocyte + Macrophage. Mouse lymph node stroma.
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
存储溶液	Preservative: 0.09% Sodium azide
纯度	Protein G purified
克隆	单克隆
克隆编号	MOMA-2
同种型	IgG2b

应用

The Abpromise guarantee Abpromise™承诺保证使用ab33451于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
IHC-Fr	★★★★★ (2)	Use at an assay dependent concentration.
IHC-P		Use at an assay dependent concentration. PubMed: 24449804
IHC-FoFr	★★★★★ (1)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent concentration. Membrane permeabilisation is required for this application. ab18536 - Rat monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

靶标

相关性

Monocyte and macrophage are white blood cells that roam the body tissues engulfing foreign organisms. A monocyte is a leukocyte, part of the human body's immune system that protects against blood-borne pathogens and moves quickly (aprox. 8-12 hours) to sites of infection in the tissues. Monocytes are usually identified in stained smears by their large bi-lobed nucleus. Macrophages are cells within the tissues that originate from specific white blood cells called monocytes. Monocytes and macrophages are phagocytes, acting in both nonspecific defense (or innate immunity) as well as specific defense (or cell-mediated immunity) of vertebrate animals. Their role is to phagocytize (engulf and then digest) cellular debris and pathogens either as stationary or mobile cells, and to stimulate lymphocytes and other immune cells to respond to the pathogen.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors