

Anti-MyD88 antibody [4D6] ab107585

敲除验证

1 Abreviews 3 References 4 图像

概述	
产品名称	Anti-MyD88抗体[4D6]
描述	小鼠单克隆抗体[4D6] to MyD88
宿主	Mouse
特异性	Expected to recognise Isoforms 1, 3, 4, 5, CRAa and CRAb.
经测试应用	适用于: Flow Cyt (Intra), WB, ICC/IF
种属反应性	与反应: Human
免疫原	Synthetic peptide, corresponding to amino acids 50-100 of Human MyD88 (NP_002459).
阳性对照	human ovary, human prostate and Jurkat cell lysates and CF-7 cells. Flow Cyt (Intra): Jurkat cells
常规说明	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>
性能	
形式	Liquid
存放说明	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
存储溶液	pH: 7.4 Preservative: 0.05% Sodium azide Constituents: PBS, 0.05% BSA
纯度	Protein G purified
克隆	单克隆
克隆编号	4D6
同种型	IgG1
轻链类型	kappa

应用

The Abpromise guarantee

Abpromise™承诺保证使用ab107585于以下的经测试应用


“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
Flow Cyt (Intra)		Use 0.5µg for 10 ⁶ cells. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
WB		Use a concentration of 5 - 7 µg/ml. Predicted molecular weight: 33 kDa.
ICC/IF		Use a concentration of 5 µg/ml.

靶标

功能	Adapter protein involved in the Toll-like receptor and IL-1 receptor signaling pathway in the innate immune response. Acts via IRAK1, IRAK2, IRF7 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Increases IL-8 transcription. Involved in IL-18-mediated signaling pathway.
组织特异性	Ubiquitous.
疾病相关	Defects in MYD88 are the cause of MYD88 deficiency (MYD88D) [MIM:612260]; also known as recurrent pyogenic bacterial infections due to MYD88 deficiency. Patients suffer from autosomal recessive, life-threatening, often recurrent pyogenic bacterial infections, including invasive pneumococcal disease, and die between 1 and 11 months of age. Surviving patients are otherwise healthy, with normal resistance to other microbes, and their clinical status improved with age.
序列相似性	Contains 1 death domain. Contains 1 TIR domain.
结构域	The intermediate domain (ID) is required for the phosphorylation and activation of IRAK.
细胞定位	Cytoplasm.

图片



Western blot - Anti-MyD88 antibody [4D6]
(ab107585)

Lane 1: Wild-type HAP1 cell lysate (20 µg)


Lane 2: MyD88 knockout HAP1 cell lysate (20 µg)

Lane 3: Jurkat cell lysate (20 µg)

Lane 4: MOLT4 cell lysate (20 µg)


Lanes 1 - 4: Merged signal (red and green). Green - ab107585 observed at 37 kDa. Red - loading control, **ab176560**, observed at 52 kDa.

ab107585 was shown to recognize MyD88 when MyD88 knockout samples were used, along with additional cross-reactive bands. Wild-type and MyD88 knockout samples were subjected to SDS-PAGE. ab107585 and **ab176560** (loading control to alpha tubulin) were diluted 5 µg/mL and 1/10000 respectively and incubated overnight at 4°C. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preadsorbed **ab216772** and Goat Anti-Rabbit IgG H&L (IRDye® 680RD) preadsorbed **ab216777** secondary antibodies at 1/10000 dilution for 1 hour at room temperature before imaging.




Flow Cytometry (Intracellular) - Anti-MyD88 antibody
[4D6] (ab107585)

Intracellular flow cytometric analysis of MyD88 antibody (ab107585) in Jurkat cells (0.5 ug). Shaded histogram represents cells without antibody; green represents isotype control; red represents anti-MyD88 antibody. Secondary antibody- anti-mouse IgG PE conjugated.



Immunocytochemistry/ Immunofluorescence - Anti-MyD88 antibody [4D6] (ab107585)

Immunocytochemical analysis of MyD88 antibody (ab107585) in MCF-7 cells using an isotype control (top) and Myd88 Antibody (5 µg/ml; bottom).



Western blot - Anti-MyD88 antibody [4D6] (ab107585)

All lanes : Anti-MyD88 antibody [4D6] (ab107585) at 5 µg/ml

Lane 1 : Human ovary cell lysate

Lane 2 : Human prostate cell lysate

Lane 3 : Jurkat cell lysate

Lane 4 : Jurkat cell lysate with Immunizing peptide

Secondary

All lanes : Goat anti-mouse Ig HRP

Predicted band size: 33 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors