

Anti-Myelin Basic Protein antibody [12] ab7349

重组

★★★★★ **21 Abreviews** **264 References** **5 图像**

概述

产品名称	Anti-Myelin Basic蛋白抗体[12]
描述	大鼠单克隆抗体[12] to Myelin Basic蛋白
宿主	Rat
经测试应用	适用于: ICC/IF, WB, IHC-P
种属反应性	与反应: Mouse, Rat, Human
免疫原	Full length protein corresponding to Cow Myelin Basic Protein.
表位	Amino acids 82-87 (DENPVV).
阳性对照	WB: Human cerebellum, Mouse brain tissue and Rat brain tissue. IHC-P: Human cerebellum tissue. ICC/IF: Mouse primary oligodendroglia cells.
常规说明	<p>This product has switched from a hybridoma to recombinant production method on 08th November 2021.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p>

性能

形式	Liquid
存放说明	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle.
存储溶液	Preservative: 0.01% Sodium azide Constituents: 59% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
纯度	Ion Exchange Chromatography
克隆	单克隆
克隆编号	12
同种型	IgG2a

应用


The Abpromise guarantee **Abpromise™**承诺保证使用ab7349于以下的经测试应用

“应用说明”部分 下显示的仅为推荐的起始稀释度;实际最佳的稀释度/浓度应由使用者检定。

应用	Ab评论	说明
ICC/IF		1/200.
WB	★★★★★ (2)	1/1000.
IHC-P	★★★★★ (6)	1/4000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

靶标

功能	The classic group of MBP isoforms (isoform 4-isoform 14) are with PLP the most abundant protein components of the myelin membrane in the CNS. They have a role in both its formation and stabilization. The smaller isoforms might have an important role in remyelination of denuded axons in multiple sclerosis. The non-classic group of MBP isoforms (isoform 1-isoform 3/Golli-MBPs) may preferentially have a role in the early developing brain long before myelination, maybe as components of transcriptional complexes, and may also be involved in signaling pathways in T-cells and neural cells. Differential splicing events combined with optional post-translational modifications give a wide spectrum of isomers, with each of them potentially having a specialized function. Induces T-cell proliferation.
组织特异性	MBP isoforms are found in both the central and the peripheral nervous system, whereas Golli-MBP isoforms are expressed in fetal thymus, spleen and spinal cord, as well as in cell lines derived from the immune system.
疾病相关	Note=The reduction in the surface charge of citrullinated and/or methylated MBP could result in a weakened attachment to the myelin membrane. This mechanism could be operative in demyelinating diseases such as chronic multiple sclerosis (MS), and fulminating MS (Marburg disease).
序列相似性	Belongs to the myelin basic protein family.
发展阶段	Expression begins abruptly in 14-16 week old fetuses. Even smaller isoforms seem to be produced during embryogenesis; some of these persisting in the adult. Isoform 4 expression is more evident at 16 weeks and its relative proportion declines thereafter.
翻译后修饰	Several charge isomers of MBP; C1 (the most cationic, least modified, and most abundant form), C2, C3, C4, C5, C6, C7, C8-A and C8-B (the least cationic form); are produced as a result of optional PTM, such as phosphorylation, deamidation of glutamine or asparagine, arginine citrullination and methylation. C8-A and C8-B contain each two mass isoforms termed C8-A(H), C8-A(L), C8-B(H) and C8-B(L), (H) standing for higher and (L) for lower molecular weight. C3, C4 and C5 are phosphorylated. The ratio of methylated arginine residues decreases during aging, making the protein more cationic. The N-terminal alanine is acetylated (isoform 3, isoform 4, isoform 5 and isoform 6). Arg-241 was found to be 6% monomethylated and 60% symmetrically dimethylated.
细胞定位	Myelin membrane. Cytoplasmic side of myelin.



Immunocytochemistry/ Immunofluorescence - Anti-Myelin Basic Protein antibody [12] (ab7349)

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized mouse primary neural/glia cells labelling Myelin Basic Protein with ab7349 at a 1/200 dilution, followed by Goat anti-Rat secondary (Alexa Fluor® 488) ([ab150157](#)) secondary antibody at a 1/1000 dilution. Confocal image showing positive staining in mouse primary oligodendroglia cells (shown in green). Nuclear DNA was labeled with DAPI (shown in blue).


Counterstained with Anti-MAP2 antibody ([ab183830](#)) at a 1/1000 dilution, followed by Goat Anti-Rabbit secondary (Alexa Fluor® 594) ([ab150080](#)) at a 1/1000 dilution (shown in magenta).

Confocal scanning Z step was set as 0.3 µm followed by image processing with maximum Z projection. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

The negative controls are as follows:

-ve control 1: Myelin Basic Protein (ab7349) at a 1/200 dilution, followed by Goat anti-Rabbit secondary (Alexa Fluor® 594) ([ab150080](#)) at a 1/1000 dilution.

-ve control 2: Anti-MAP2 ([ab183830](#)) at a 1/1000 dilution, followed by Goat anti-Rat secondary (Alexa Fluor® 488) ([ab150157](#)) at a 1/1000 dilution.



Western blot - Anti-Myelin Basic Protein antibody [12] (ab7349)

Anti-Myelin Basic Protein antibody [12] (ab7349) at 1/5000 dilution + Human cerebellum tissue lysate at 10 µg

Secondary

Goat Anti-Rat IgG H&L (HRP) ([ab205720](#)) at 1/10000 dilution


Developed using the ECL technique.

Observed band size: 18.5 kDa

Exposure time: 1 second

The molecular weight observed was in consistency with the literature (PMID: 9299539).

Blocking buffer: 5% NFDM/TBST.



Western blot - Anti-Myelin Basic Protein antibody
[12] (ab7349)

All lanes : Anti-Myelin Basic Protein antibody [12] (ab7349) at
1/5000 dilution

Lane 1 : Mouse brain tissue lysate

Lane 2 : Rat brain tissue lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat Anti-Rat IgG H&L (HRP) (**ab205720**) at 1/10000
dilution

Observed band size: 14-21.5 kDa

The molecular weight observed was in consistency with the
literature (PMID: 9299539).

Blocking buffer: 5% NFDm/TBST.


Exposure time: Lane 1: 3 seconds; Lane 2: 1 second.



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Myelin Basic Protein
antibody [12] (ab7349)

Immunohistochemical analysis of paraffin-embedded Human
cerebrum labeling Myelin Basic Protein with ab7349 at 1/4000
dilution (0.259 µg/ml) followed by a ready to use Goat Anti-Rat IgG
H&L (HRP polymer; **ab214882**). The section was incubated with
ab7349 for 30 mins at room temperature. The immunostaining was
performed on a Leica Biosystems BOND® RX instrument.
Counterstained with Hematoxylin.

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope
retrieval solution 1) for 20 mins.



Immunohistochemical analysis of paraffin-embedded Human cerebrum labeling Myelin Basic Protein with ab7349 at 1/4000 dilution (0.259 µg/ml) followed by a ready to use Goat Anti-Rat IgG H&L (HRP polymer; **ab214882**). The section was incubated with ab7349 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with DAPI.

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.cn/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors